25





Novel polynucleotides and polypeptides in pathogenic mycobacteria and their use as diagnostics, vaccines and targets for chemotherapy.

This invention relates to the novel polynucleotide sequence we have designated "GS" which we have identified in pathogenic mycobacteria. GS is a pathogenicity island within 8kb of DNA comprising a core region of 5.75kb and an adjacent transmissable element within 2.25kb. GS is contained within Mycobacterium paratuberculosis, Mycobacterium avium subsp. silvaticum and some pathogenic isolates of M.avium. Functional portions of the core region of GS are also represented by regions with a high degree of homology that we have identified in cosmids containing genomic DNA from Mycobacterium tuberculosis.

15 Background to the invention

Mycobacterium tuberculosis (Mtb) is a major cause of global diseases of humans as well as animals. Although conventional methods of diagnosis including microscopy, culture and skin testing exist for the recognition of these diseases, improved methods particularly new immunodiagnostics and DNA-based detection systems are needed. Drugs used to treat tuberculosis are increasingly encountering the problem of resistant organisms. New drugs targeted at specific pathogenicity determinants as well as new vaccines for the prevention and treatment of tuberculosis are required. The importance of Mtb as a global pathogen is reflected in the commitment being made to sequencing the entire genome of this organism. This has generated a large amount of DNA sequence data of genomic DNA within cosmid and other libraries. Although the DNA sequence is known in the art, the functions of the vast majority of these sequences, the proteins they encode, the biological significance of these proteins, and the overall relevance and use of these genes and their products as diagnostics, vaccines and targets for chemotherapy for tuberculous disease, remains entirely unknown.

35 Mycobacterium avium subsp.silvaticum (Mavs) is a pathogenic mycobacterium causing diseases of animals and birds, but it can

35



also affect humans. Mycobacterium paratuberculosis (Mptb) causes chronic inflammation of the intestine in many species of animals including primates and can also cause Crohn's disease in humans. Mptb is associated with other chronic inflammatory diseases of Subclinical Mptb infection is humans such as sarcoidosis. widespread in domestic livestock and is present in milk from resistant more is The organism infected animals. pasteurisation than Mtb and can be conveyed to humans in retail Mptb is also present in water supplies, milk supplies. particularly those contaminated with run-off from heavily grazed pastures. Mptb and Mavs contain the insertion elements IS900 and IS902 respectively, and these are linked to pathogenicity in IS900 and IS902 provide convenient highly these organisms. specific multi-copy DNA targets for the sensitive detection of these organisms using DNA-based methods and for the diagnosis of infections in animals and humans. Much improvement is however required in the immunodiagnosis of Mptb and Mavs infections in animals and humans. Mptb and Mavs are in general, resistant in vivo to standard anti-tuberculous drugs. Although substantial clinical improvements in infections caused by Mptb, such as may result from treatment of patients with Crohn's disease, combinations of existing drugs such as Rifabutin, Clarithromycin additional effective drug treatments are or Azithromycin, Furthermore, there is an urgent need for effective required. vaccines for the prevention and treatment of Mptb and Mavs 25 infections in animals and humans based upon the recognition of specific pathogenicity determinants.

Pathogenicity islands are, in general, 7-9kb regions of DNA comprising a core domain with multiple ORFs and an adjacent transmissable element. The transmissable element also encodes proteins which may be linked to pathogenicity, such as by providing receptors for cellular recognition. Pathogenicity islands are envisaged as mobile packages of DNA which, when they enter an organism, assist in bringing about its convertion from a non-disease-causing to a disease-causing strain.

Description of the Drawings

20

25

Figure 1(a) and (b) shows a linear map of the pathogenicity island GS in Mavs (Fig 1a) and in Mptb (Fig 1b). The main open reading frames are illustrated as ORFs A to H. ORFs A to F are found within the core region of GS. ORFs G and H are encoded by the adjacent transmissable element portion of GS.

Disclosure of the invention

Using a DNA-based differential analysis technology we have discovered and characterised a novel polynucleotide in Mptb (isolates 0022 from a Guernsey cow and 0021 from a red deer). This polynucleotide comprises the gene region we have designated GS is found in Mptb using the identifier DNA sequences Seq.ID.No 1 and 2 where the Seq.ID No2 is the complementary sequence of Seq.ID No 1. GS is also identified in Mavs. complete DNA sequence incorporating the positive strand of GS from an isolate of Mavs comprising 7995 nucleotides, including the core region of GS and adjacent transsmissable element, is given in Seq.ID No.3. DNA sequence comprising 4435 bp of the positive strand of GS obtained from an isolate of Mptb including the core region of GS (nucleotides 1614 to 6047 of GS in Mavs) is given in Seq.ID No 4. The DNA sequence of GS from Mptb is highly (99.4%) homologous to GS in Mavs. The remaining portion of the DNA sequence of GS in Mptb, is readily obtainable by a person skilled in the art using standard laboratory procedures. The entire functional DNA sequence including core region and transmisable element of GS in Mptb and Mavs as described above, comprise the polynucleotide sequences of the invention.

There are 8 open reading frames (ORFs) in GS. Six of these designated GSA, GSB, GSC, GSD, GSE and GSF are encoded by the core DNA region of GS which, characteristically for a pathogenicity island, has a different GC content than the rest of the microbial genome. Two ORFs designated GSG and GSH are encoded by the transmissable element of GS whose GC content resembles that of the rest of the mycobacterial genome. The ORF GSH comprises two sub-ORFs H₁ H₂ on the complementary DNA strand linked by a programmed frameshifting site so that a single polypeptide is translated from the ORF GSH. The nucleotide

20

25

15.

sequences of the 8 ORFs in GS and their translations are shown in Seq. ID No 5 to Seq.ID No 29 as follows:

ORF A: Seq. ID No 5 Nucleotides 50 to 427 of GS from Mavs Seq. ID No 6 Amino acid sequence encoded by Seq. ID No 5.

ORF B: Seq. ID No 7 Nucleotides 772 to 1605 of GS from Mavs Seq. ID No 8 Amino acid sequence encoded by Seq. ID No 7.

ORF C: Seq. ID No 9 Nucleotides 1814 to 2845 of GS from Mavs

Seq. ID No 10 Amino acid sequence encoded by Seq.ID No

9.

Seq. ID No 11 Nucleotides 201 to 1232 of GS from Mptb

Seq. ID No 12 Amino acid sequence encoded by Seq.ID No

11

15 ORF D: Seq. ID No 13 Nucleotides 2785 to 3804 of GS from Mavs
Seq. ID No 14 Amino acid sequence encoded by Seq. ID No
13.
Seq. ID No 15 Nucleotides 1172 to 2191 of GS from Mptb
Seq. ID No 16 Amino acid sequence encoded by Seq. ID No

ORF E: Seq. ID No 17 Nucleotides 4080 to 4802 of GS from Mavs
Seq. ID No 18 Amino acid sequence encoded by Seq. ID No
17.
Seq. ID No 19 Nucleotides 2467 to 3189 of GS from Mptb

Seq. ID No 19 Nucleotides 2407 to 3109 of GS 110 M Mpts Seq. ID No 20 Amino acid sequence encoded by Seq. ID No 19.

ORF F: Seq. ID No 21 Nucleotides 4947 to 5747 of GS from Mavs
Seq. ID No 22 Amino acid sequence encoded by Seq. ID No
21.

Seq. ID No 23 Nucleotides 3335 to 4135 of GS from Mpth Seq. ID No 24 Amino acid sequence encoded by Seq.ID No 23.



ORF G: Seq. ID No 25 Nucleotides 6176 to 7042 of GS from Mavs Seq. ID No 26 Amino acid sequence encoded by Seq. ID No 25.

ORF H: Seq.ID No 27 Nucleotides 7953 to 6215 from Mavs.

5 ORF H_1 : Seq.ID No 28 Amino acid sequence encoded by nucleotides 7953 to 7006 of Seq.ID No 27

ORF H₂: Seq.ID No 29 Amino acid sequence encoded by nucleotides 7009 to 6215 of Seq.ID No 27

The polynucleotides in Mtb with homology to the ORFs B, C, E and 10 F of GS in Mptb and Mavs, and the polypeptides they are now known to encode as a result of our invention, are as follows:

ORF B: Seq.ID No 30 Cosmid MTCY277 nucleotides 35493 to 34705

Seq.ID No 31 Amino acid sequence encoded by Seq.ID No30.

ORF C: Seq.ID No 32 Cosmid MTCY277 nucleotides 31972 to 32994 Seq.ID No 33 Amino acid sequence encoded by Seq.ID No32.

ORF E: Seq.ID No 34 Cosmid MTCY277 nucleotides 34687 to 33956

Seq.ID No 35 Amino acid sequence encoded by Seq.ID

No34.

ORF E: Seq.ID No 36 Cosmid MTO24 nucleotides 15934 to 15203 Seq.ID No 37 Amino acid sequence encoded by Seq.ID No36.

25 ORF F: Seq.ID No38 Cosmid MTO24 nucleotides 15133 to 14306 Seq.ID No 39 Amino acid sequence encoded by Seq.ID No38.

The proteins and peptides encoded by the ORFs A to H in Mptb and Mavs and the amino acid sequences from homologous genes we have

obtained.

discovered in Mtb given in Seq.ID Nos 31, 33, 35, 37 and 39, as described above and fragments thereof, comprise the polypeptides of the invention. The polypeptides of the invention are believed to be associated with specific immunoreactivity and with the pathogenicity of the host micro-organisms from which they were

The present invention thus provides a polynucleotide in substantially isolated form which is capable of selectively hybridising to sequence ID Nos 3 or 4 or a fragment thereof. The polynucleotide fragment may alternatively comprise a sequence selected from the group of Seq.ID.No: 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25 and 27. The invention further provides a polynucleotide in substantially isolated form whose sequence consists essentially of a sequence selected from the group Seq ID Nos. 30, 32, 34, 36 and 38, or a corresponding sequence selectively hybridizable thereto, or a fragment of said sequence or corresponding sequence.

The invention further provides diagnostic probes such as a probe which comprises a fragment of at least 15 nucleotides of a polynucleotide of the invention, or a peptide nucleic acid or similar synthetic sequence specific ligand, optionally carrying a revealing label. The invention also provides a vector carrying a polynucleotide as defined above, particularly an expression vector.

25 The invention further provides a polypeptide in substantially isolated form which comprises any one of the sequences selected from the group consisting Seq.ID.No: 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 29, 31, 33, 35, 37 and 39, or a polypeptide substantially homologous thereto. The invention additionally provides a polypeptide fragment which comprises a fragment of a polypeptide defined above, said fragment comprising at least 10 amino acids and an epitope. The invention also provides polynucleotides in substantially isolated form which encode polypeptides of the invention, and vectors which comprise such polynucleotides, as well as antibodies capable of binding such polypeptides. In an additional aspect, the invention provides

· D

4

20

25

30

kits comprising polynucleotides, polypeptides, antibodies or synthetic ligands of the invention and methods of using such kits in diagnosing the presence or absence of mycobacteria in a sample. The invention also provides pharmaceutical compositions comprising polynucleotides of the invention, polypeptides of the invention or antisense probes and the use of such compositions the treatment or prevention of diseases caused by The invention also provides polynucleotihe mycobacteria. prevention and treatment of infections due to GS-containing pathogenic mycobacteria in animals and humans and as a means of enhacing in vivo susceptibility of said mycobacteria to The invention also provides bacteria or antimicrobial drugs. viruses transformed with polynucleotides of the invention for use as vaccines. The invention further provides Mptb or Mavs which all or part or the polynucleotides of the invention have been deleted or disabled to provide mutated organisms of lower pathogenicity for use as vaccines in animals and humans. invention further provides Mtb in which all or part of the polynucleotides encoding polypeptides of the invention have been deleted or disabled to provided mutated organisms or lower pathogenicity for use as vaccines in animals and humans.

A further aspect of the invention is our discovery of homologies between the ORFs B, C and E in GS on the one hand, and Mtb cosmid MTCY277 on the other (data from Genbank database using the tomputer programmes BLAST and BLIXEM). The homologous ORFs in MTCY277 are adjacent to one another consistent with the form of another pathogenicity island in Mtb. A further aspect of the invention is our discovery of homologies between ORFs E and F in GS, and Mtb cosmid MTO24 (also Genbank, as above) with the homologous ORFs close to one another. The use of polynucleotides and polypeptides from Mtb (Seq. ID Nos 30,31, 32, 33, 34, 35, 36, 37, 38 and 39) in substantially isolated form as diagnostics, vaccines and targets for chemotherapy, for the management and prevention of Mtb infections in humans and animals, and the processes involved in the preparation and use of diagnostics, vaccines and new chemotherapeutic agents, comprise further aspects of the invention.

20

30

-8-

Detailed description of the invention.

A. Polynucleotides

Polynucleotides of the invention as defined herein may comprise DNA or RNA. They may also be polynucleotides which include within them synthetic or modified nucleotides or peptide nucleic A number of different types of modification to in the art. These include oligonucleotides are known methylphosphonate and phosphorothioate backbones, addition of acridine or polylysine chains at the 3' and/or 5' ends of the molecule. For the purposes of the present invention, it is to be understood that the polynucleotides described herein may be modified by any method available in the art. Such modifications may be carried out in order to couple the said polynucleotide to a solid phase or to enhance the recognition, the in vivo activity, or the lifespan of polynucleotides of the invention.

A number of different types of polynucleotides of the invention In the broadest aspect, polynucleotides and are envisaged. fragments thereof capable of hybridizing to SEQ ID NO:3 or 4 form first aspect of the invention. includes This Within this class of polynucleotide of SEQ ID NO: 3 or 4. polynucleotides various sub-classes of polynucleotides are of particular interest.

One sub-class of polynucleotides which is of interest is the class of polynucleotides encoding the open reading frames A, B, C, D, E, F, G and H, including SEQ ID NOs:5, 7, 9, 11, 13, 15, 25 17, 19, 21, 23, 25 and 27. As discussed below, polynucleotides encoding ORF H include the polynucleotide sequences 7953 to 7006 and 7009 to 6215 within SEQ ID NO: 27, as well as modified sequences in which the frame-shift has been modified so that the two sub-reading frames are placed in a single reading frame. This may be desirable where the polypeptide is to be produced in recombinant expression systems.

The invention thus provides a polynucleotide in substantially isolated form which encodes any one of these ORFs or combinations

25



thereof. Combinations thereof includes combinations of 2, 3, 4, Polynucleotides may be provided which 5 or all of the ORFs. comprise an individual ORF carried in a recombinant vector including the vectors described herein. Thus in one preferred 5 aspect the invention provides a polynucleotide in substantially isolated form capable of selectively hybridizing to the nucleic acid comprising ORFs A to F of the core region of the Mptb and Mavs pathogenicity islands of the invention. Fragments thereof corresponding to ORFs A to E, B to F, A to D, B to E, A to C, B to D or any two adjacent ORFs are also included in the invention.

Polynucleotides of the invention will be capable of selectively hybridizing to the corresponding portion of the GS region, or to the corresponding ORFs of Mtb described herein. The term "selectively hybridizing" indicates that the polynucleotides will hybridize, under conditions of medium to high stringency (for example 0.03 M sodium chloride and 0.03 M sodium citrate at from about 50°C to about 60°C) to the corresponding portion of SEQ ID NO:3 or 4 or the complementary strands thereof but not to genomic DNA from mycobacteria which are usually non-pathogenic including Such polynucleotides will non-pathogenic species of M.avium. 20 generally be generally at least 68%, e.g. at least 70%, preferably at least 80 or 90% and more preferably at least 95% homologous to the corresponding DNA of GS. The corresponding portion will be of over a region of at least 20, preferably at least 30, for instance at least 40, 60 or 100 or more contiguous nucleotides.

By "corresponding portion" it is meant a sequence from the GS region of the same or substantially similar size which has been determined, for example by computer alignment, to have the greatest degree of homology to the polynucleotide.

Any combination of the above mentioned degrees of homology and minimum sizes may be used to define polynucleotides of the invention, with the more stringent combinations (i.e. higher homology over longer lengths) being preferred. Thus for example a polynucleotide which is at least 80% homologous over 25, preferably 30 nucleotides forms one aspect of the invention, as

30

does a polynucleotide which is at least 90% homologous over 40 nucleotides.

A further class of polynucleotides of the invention is the class of polynucleotides encoding polypeptides of the invention, the 5 polypeptides of the invention being defined in section B below. the redundancy of the genetic code polynucleotides may be of a lower degree of homology than required for selective hybridization to the GS region. However, when such polynucleotides encode polypeptides of the invention these polynucleotides form a further aspect. It may for example be desirable where polypeptides of the invention are produced recombinantly to increase the GC content of such polynucleotides. This increase in GC content may result in higher levels of expression via codon usage more appropriate to the host cell in which recombinant expression is taking place.

An additional class of polynucleotides of the invention are those obtainable from cosmids MTCY277 and MT024 (containing Mtb genomic sequences), which polynucleotides consist essentially of the fragment of the cosmid containing an open reading frame encoding any one of the homologous ORFs B, C, E or F respectively. polynucleotides are referred to below as Mtb polynucleotides. However, where reference is made to polynucleotides in general such reference includes Mtb polynucleotides unless the context In addition, the invention is explicitly to the contrary. provides polynucleotides which encode the same polypeptide as the abovementioned ORFs of Mtb but which, due to the redundancy of the genetic code, have different nucleotide sequences. form further Mtb polynucleotides of the invention. Fragments of Mtb polynucleotides suitable for use as probes or primers also form a further aspect of the invention.

The invention further provides polynucleotides in substantially isolated form capable of selectively hybridizing (where selectively hybridizing is as defined above) to the Mtb polynucleotides of the invention.



The invention further provides the Mtb polynucleotides of the invention linked, at either the 5' and/or 3' end to polynucleotide sequences to which they are not naturally contiguous. Such sequences will typically be sequences found in cloning or expression vectors, such as promoters, 5' untranslated sequence, 3' untranslated sequence or termination sequences. The sequences may also include further coding sequences such as signal sequences used in recombinant production of proteins.

Further polynucleotides of the invention are illustrated in the accompanying examples.

Polynucleotides of the invention may be used to produce a primer, e.g. a PCR primer, a primer for an alternative amplification reaction, a probe e.g. labelled with a revealing label by conventional means using radioactive or non-radioactive labels or a probe linked covalently to a solid phase, or the polynucleotides may be cloned into vectors. Such primers, probes and other fragments will be at least 15, preferably at least 20, for example at least 25, 30 or 40 or more nucleotides in length, and are also encompassed by the term polynucleotides of the invention as used herein.

Primers of the invention which are preferred include primers directed to any part of the ORFs defined herein. The ORFs from other isolates of pathogenic mycobacteria which contain a GS. region may be determined and conserved regions within each Primers directed to such individual ORF may be identified. conserved regions form a further preferred aspect of the In addition, the primers and other polynucleotides of the invention may be used to identify, obtain and isolate ORFs capable of selectively hybridizing to the polynucleotides of the invention which are present in pathogenic mycobacteria but which are not part of a pathogenicity island in that particular species Thus in addition to the ORFs B, C, E and F which of bacteria. have been identified in Mtb, similar ORFs may be identified in other pathogens and ORFs corresponding to the GS ORFs C, D, E, F and H, may also be identified.

Polynucleotides such as DNA polynucleotides and probes according to the invention may be produced recombinantly, synthetically, or by any means available to those of skill in the art. They may also be cloned by standard techniques.

In general, primers will be produced by synthetic means, involving a step-wise manufacture of the desired nucleic acid sequence one nucleotide at a time. Techniques for accomplishing this using automated techniques are readily available in the art. Longer polynucleotides will generally be produced using recombinant means, for example using a PCR (polymerase chain reaction) cloning techniques. This will involve making a pair or primers (e.g. of about 15-30 nucleotides) to a region of GS, which it is desired to clone, bringing the primers into contact with genomic DNA from a mycobacterium or a vector carrying the GS sequence, performing a polymerase chain reaction under 15 conditions which bring about amplification of the desired region, isolating the amplified fragment (e.g. by purifying the reaction mixture on an agarose gel) and recovering the amplified DNA. The primers may be designed to contain suitable restriction enzyme recognition sites so that the amplified DNA can be cloned into a suitable cloning vector.

Such techniques may be used to obtain all or part of the GS or ORF sequences described herein, as well as further genomic clones containing full open reading frames. Although in general such techniques are well known in the art, reference may be made in particular to Sambrook J., Fritsch EF., Maniatis T (1989). Molecular cloning: a Laboratory Manual, 2nd edn. Cold Spring Harbor, New York, Cold Spring Harbor Laboratory.

Polynucleotides which are not 100% homologous to the sequences of the present invention but fall within the scope of the invention can be obtained in a number of ways.

Other isolates or strains of pathogenic mycobacteria will be expected to contain allelic variants of the GS sequences described herein, and these may be obtained for example by probing genomic DNA libraries made from such isolates or strains

of bacteria using GS or ORF sequences as probes under conditions of medium to high stringency (for example 0.03M sodium chloride and 0.03M sodium citrate at from about 50°C to about 60°C).

A particularly preferred group of pathogenic mycobacteria are isolates of *M.paratuberculosis*. Polynucleotides based on GS regions from such bacteria are particularly preferred. Preferred fragments of such regions include fragments encoding individual open reading frames including the preferred groups and combinations of open reading frames discussed above.

Alternatively, such polynucleotides may be obtained by site directed mutagenesis of the GS or ORF sequences or allelic variants thereof. This may be useful where for example silent codon changes are required to sequences to optimise codon which for a particular host cell in polynucleotide sequences are being expressed. Other sequence changes may be desired in order to introduce restriction enzyme recognition sites, or to alter the property or function of the polypeptides encoded by the polynucleotides of the invention. Such altered property or function will include the addition of amino acid sequences of consensus signal peptides known in the 20 art to effect transport and secretion of the modified polypeptide Another altered property will include of the invention. metagenesis of a catalytic residue or generation of fusion proteins with another polypeptide. Such fusion proteins may be with an enzyme, with an antibody or with a cytokine or other ligand for a receptor, to target a polypeptide of the invention to a specific cell type in vitro or in vivo.

The invention further provides double stranded polynucleotides comprising a polynucleotide of the invention and its complement.

Polynucleotides or primers of the invention may carry a revealing label. Suitable labels include radioisotopes such as ³²P or ³⁵S, enzyme labels, other protein labels or smaller labels such as biotin or fluorophores. Such labels may be added to polynucleotides or primers of the invention and may be detected using by techniques known per se.

15

20

25

- 14 -

Polynucleotides or primers of the invention or fragments thereof labelled or unlabelled may be used by a person skilled in the art in nucleic acid-based tests for the presence or absence of Mptb, Mavs, other GS-containing pathogenic mycobacteria, or Mtb applied to samples of body fluids, tissues, or excreta from animals and humans, as well as to food and environmental samples such as river or ground water and domestic water supplies.

Human and animal body fluids include sputum, blood, serum, plasma, saliva, milk, urine, csf, semen, faeces and infected discharges. Tissues include intestine, mouth ulcers, skin, lymph nodes, spleen, lung and liver obtained surgically or by a biopsy technique. Animals particularly include commercial livestock such as cattle, sheep, goats, deer, rabbits but wild animals and animals in zoos may also be tested.

Such tests comprise bringing a human or animal body fluid or tissue extract, or an extract of an environmental or food sample, into contact with a probe comprising a polynucleotide or primer of the invention under hybridising conditions and detecting any duplex formed between the probe and nucleic acid in the sample. Such detection may be achieved using techniques such as PCR or by immobilising the probe on a solid support, removing nucleic acid in the sample which is not hybridized to the probe, and then detecting nucleic acid which has hybridized to the probe. Alternatively, the sample nucleic acid may be immobilized on a solid support, and the amount of probe bound to such a support can be detected. Suitable assay methods of this any other formats can be found in for example WO89/03891 and WO90/13567.

Polynucleotides of the invention or fragments thereof labelled or unlabelled may also be used to identify and characterise different strains of Mptb, Mavs, other GS-containing pathogenic mycobacteria, or Mtb, and properties such as drug resistance or susceptibility.

The probes of the invention may conveniently be packaged in the form of a test kit in a suitable container. In such kits the probe may be bound to a solid support where the assay format for

25

which the kit is designed requires such binding. The kit may also contain suitable reagents for treating the sample to be probed, hybridising the probe to nucleic acid in the sample, control reagents, instructions, and the like.

The use of polynucleotides of the invention in the diagnosis of inflammatory diseases such as Crohn's disease or sarcoidosis in humans or Johne's disease in animals form a preferred aspect of the invention. The polynucleotides may also be used in the prognosis of these diseases. For example, the response of a human or animal subject in response to antibiotic, vaccination or other therapies may be monitored by utilizing the diagnostic methods of the invention over the course of a period of treatment and following such treatment.

The use of Mtb polynucleotides (particularly in the form of probes and primers) of the invention in the above-described methods form a further aspect of the invention, particularly for the detection, diagnosis or prognosis of Mtb infections.

B. Polypeptides.

polypeptides Polypeptides of invention include the substantially isolated form encoded by GS. This includes the encoded by the positive length polypeptides full complementary negative strands of GS. Each of the full length polypeptides will contain one of the amino acid sequences set out in Seq ID NOs:6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28 and Polypeptides of the invention further include variants of such sequences, including naturally occurring allelic variants and synthetic variants which are substantially homologous to said polypeptides. In this context, substantial homology is regarded as a sequence which has at least 70%, e.g. 80%, 90%, 95% or 98% amino acid homology (identity) over 30 or more, e.g 40, 50 or 100 amino acids. For example, one group of substantially homolgous polypeptides are those which have at least 95% amino acid identity to a polypeptide of any one of Seq ID NOs:6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28 and 29 over their entire length.

35 Even more preferably, this homology is 98%.

25

30

35

Polypeptides of the invention further include the polypeptide sequences of the homologous ORFs of Mtb, namely Seq ID Nos. 31, 33, 35, 37 and 39. Unless explicitly specified to the contrary, reference to polypeptides of the invention and their fragments include these Mtb polypeptides and fragments, and variants thereof (substanially homologous to said sequences) as defined herein.

Polypeptides of the invention may be obtained by the standard techniques mentioned above. Polypeptides of the invention also include fragments of the above mentioned full length polypeptides and variants thereof, including fragments of the sequences set out in SEQ ID NOs:6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 29, 31, 33, 35, 37 and 39. Such fragments for example of 8, 10, 12, 15 or up to 30 or 40 amino acids may also be obtained synthetically using standard techniques known in the art.

Preferred fragments include those which include an epitope, especially an epitope which is specific to the pathogenicity of the mycobacterial cell from which the polypeptide is derived. Suitable fragments will be at least about 5, e.g. 8, 10, 12, 15 or 20 amino acids in size, or larger. Epitopes may be determined either by techniques such as peptide scanning techniques as described by Geysen et al, Mol.Immunol., 23; 709-715 (1986), as well as other techniques known in the art.

The term "an epitope which is specific to the pathogenicity of the mycobacterial cell" means that the epitope is encoded by a portion of the GS region, or by the corresponding ORF sequences of Mtb which can be used to distinguish mycobacteria which are pathogenic by from related non-pathogenic mycobacteria including non-pathogenic species of M.avium. This may be determined using routine methodology. A candidate epitope from an ORF may be prepared and used to immunise an animal such as a rat or rabbit in order to generate antibodies. The antibodies may then be used to detect the presence of the epitope in pathogenic mycobacteria and to confirm that non-pathogenic mycobacteria do not contain any proteins which react with the epitope. Epitopes may be linear or conformational.

- 17 -

Polypeptides of the invention may be in a substantially isolated form. It will be understood that the polypeptide may be mixed with carriers or diluents which will not interfere with the intended purpose of the polypeptide and still be regarded as substantially isolated. A polypeptide of the invention may also be in a substantially purified form, in which case it will generally comprise the polypeptide in a preparation in which more than 90%, e.g. 95%, 98% or 99% of the polypeptide in the preparation is a polypeptide of the invention.

Polypeptides of the invention may be modified to confer a desired property or function for example by the addition of Histidine residues to assist their purification or by the addition of a signal sequence to promote their secretion from a cell.

Thus, polypeptides of the invention include fusion proteins which comprise a polypeptide encoding all or part of one or more of an ORF of the invention fused at the N- or C-terminus to a second sequence to provide the desired property or function. Sequences which promote secretion from a cell include, for example the yeast α -factor signal sequence.

A polypeptide of the invention may be labelled with a revealing 20 The revealing label may be any suitable label which label. allows the polypeptide to be detected. Suitable labels include radioisotopes, e.g. 125I, 35S enzymes, antibodies, polynucleotides Labelled polypeptides of the such as biotin. and ligands invention may be used in diagnostic procedures such as 25 immunoassays in order to determine the amount of a polypeptide Polypeptides or labelled of the invention in a sample. polypeptides of the invention may also be used in serological or cell mediated immune assays for the detection of immune reactivity to said polypeptides in animals and humans using 30 standard protocols.

A polypeptide or labelled polypeptide of the invention or fragment thereof may also be fixed to a solid phase, for example the surface of an immunoassay well, microparticle, dipstick or biosensor. Such labelled and/or immobilized polypeptides may be

20

30

35

packaged into kits in a suitable container along with suitable reagents, controls, instructions and the like.

Such polypeptides and kits may be used in methods of detection of antibodies or cell mediated immunoreactivity, to the mycobacterial proteins and peptides encoded by the ORFs of the invention and their allelic variants and fragments, using immunoassay. Such host antibodies or cell mediated immune reactivity will occur in humans or animals with an immune system which detects and reacts against polypeptides of the invention.

The antibodies may be present in a biological sample from such humans or animals, where the biological sample may be a sample as defined above particularly blood, milk or saliva.

Immunoassay methods are well known in the art and will generally comprise:

(a) providing a polypeptide of the invention comprising an epitope bindable by an antibody against said mycobacterial polypeptide;

(b) incubating a biological sample with said polypeptide under conditions which allow for the formation of an antibody-antigen complex; and

(c) determining whether antibody-antigen complex comprising said polypeptide is formed.

Immunoassay methods for cell mediated immune reactivity in animals and humans are also well known in the art (e.g. as described by Weir et al 1994, J.Immunol Methods 176; 93-101) and will generally comprise

- (a) providing a polypeptide of the invention comprising an epitope bindable by a lymphocyte or macrophage or other cell receptor;
- (b) incubating a cell sample with said polypeptide under conditions which allow for a cellular immune response such as release of cytokines or other mediator to occur; and
- (c) detecting the presence of said cytokine or mediator in the incubate.

20

25

Polypeptides of the invention may be made by standard synthetic means well known in the art or recombinantly, as described below.

Polypeptides of the invention or fragments thereof labelled or unlabelled may also be used to identify and characterise different strains of Mptb, Mavs, other GS-containing pathogenic mycobacteria, or Mtb, and properties such as drug resistance or susceptibility.

The polypeptides of the invention may conveniently be packaged in the form of a test kit in a suitable container. In such kits the polypeptide may be bound to a solid support where the assay format for which the kit is designed requires such binding. The kit may also contain suitable reagents for treating the sample to be examined, control reagents, instructions, and the like.

The use of polypeptides of the invention in the diagnosis of inflammatory diseases such as Crohn's disease or sarcoidosis in humans or Johne's disease in animals form a preferred aspect of the invention. The polypeptides may also be used in the prognosis of these diseases. For example, the response of a human or animal subject in response to antibiotic or other therapies may be monitored by utilizing the diagnostic methods of the invention over the course of a period of treatment and following such treatment.

The use of Mtb polypeptides of the invention in the above-described methods form a further aspect of the invention, particularly for the detection, diagnosis or prognosis of Mtb infections.

Polypeptides of the invention may also be used in assay methods for identifying candidate chemical compounds which will be useful in inhibiting, binding to or disrupting the function of said polypeptides required for pathogenicity. In general, such assays involve bringing the polypeptide into contact with a candidate inhibitor compound and observing the ability of the compound to disrupt, bind to or interfer with the polypeptide.

25

30

35

.

There are a number of ways in which the assay may be formatted. For example, those polypeptides which have an enzymatic function may be assayed using labelled substrates for the enzyme, and the amount of, or rate of, conversion of the substrate into a product measured, e.g by chromatograpy such as HPLC or by a colourimetric assay. Suitable labels include ³⁵S, ¹²⁵I, biotin or enzymes such as horse radish peroxidase.

For example, the gene product of ORF C is believed to have GDP-mannose dehydratase activity. Thus an assay for inhbitors of the gene product may utilise for example labelled GDP-mannose, GDP or mannose and the activity of the gene product followed. ORF D encodes a gene related to the synthesis and regulation of capuslar polysaccharides, which are often associated with invasiveness and pathogenicity. Labelled polysaccharide substrates may be used in assays of the ORF D gene product. The gene product of ORF F encodes a protein with putative glucosyl transferase activity and thus labelled amino sugars such as $\beta\textsc{-1}\textsc{-3}\textsc{-N-acetylglucosamine}$ may be used as substrates in assays.

Candidate chemical compounds which may be used may be natural or synthetic chemical compounds used in drug screening programmes. Extracts of plants which contain several characterised or uncharacterised components may also be used.

Alternatively, the a polypeptide of the invention may be screened against a panel of peptides, nucleic acids or other chemical functionalities which are generated by combinatorial chemistry. This will allow the definition of chemical entities which bind to polypeptides of the invention. Typically, the polypeptide of the invention will be brought into contact with a panel of compounds from a combinantorial library, with either the panel or the polypeptide being immobilized on a solid phase, under conditions suitable for the polypeptide to bind to the panel. The solid phase will then be washed under conditions in which only specific interactions between the polypeptide and individual members of the panel are retained, and those specific members may be utilized in further assays or used to design further panels of candidate ompounds.

30

For example, a number of assay methods to define peptide interaction with peptides are known. For example, W086/00991 describes a method for determining mimotopes which comprises making panels of catamer preparations, for example octamers of amino acids, at which one or more of the positions is defined and the remaining positions are randomly made up of other amino acids, determining which catamer binds to a protein of interest and re-screening the protein of interest against a further panel based on the most reactive catamer in which one or more additional designated positions are systematically varied. This may be repeated throughout a number of cycles and used to build up a sequence of a binding candidate compound of interest.

WO89/03430 describes screening methods which permit the preparation of specific mimotopes which mimic the immunological activity of a desired analyte. These mimotopes are identified by reacting a panel of individual peptides wherein said peptides are of systematically varying hydrophobicity, amphipathic characteristics and charge patterns, using an antibody against an antigen of interest. Thus in the present case antibodies against the a polypeptide of the inventoin may be employed and mimotope peptides from such panels may be identified.

C. Vectors.

Polynucleotides of the invention can be incorporated into a recombinant replicable vector. The vector may be used to replicate the nucleic acid in a compatible host cell. Thus in a further embodiment, the invention provides a method of making polynucleotides of the invention by introducing a polynucleotide of the invention into a replicable vector, introducing the vector into a compatible host cell, and growing the host cell under conditions which bring about replication of the vector. The vector may be recovered from the host cell. Suitable host cells are described below in connection with expression vectors.

D. Expression Vectors.

15

20

25

35

Preferably, a polynucleotide of the invention in a vector is operably linked to a control sequence which is capable of providing for the expression of the coding sequence by the host cell, i.e. the vector is an expression vector. The term "operably linked" refers to a juxtaposition wherein the components described are in a relationship permitting them to function in their intended manner. A control sequence "operably linked" to a coding sequence is ligated in such a way that expression of the coding sequence is achieved under conditions compatible with the Such vectors may be transformed into a control sequences. suitable host cell as described above to provide for expression of a polypeptide of the invention. Thus, in a further aspect the invention provides a process for preparing polypeptides according to the invention which comprises cultivating a host transformed or transfected with an expression vector as described above, under conditions to provide for expression by the vector of a coding sequence encoding the polypeptides, and recovering the expressed polypeptides.

A further embodiment of the invention provides vectors for the replication and expression of polynucleotides of the invention, or fragments thereof. The vectors may be for example, plasmid, virus or phage vectors provided with an origin of replication, optionally a promoter for the expression of polynucleotide and optionally a regulator of the promoter. The vectors may contain one or more selectable marker genes, for example an ampicillin resistance gene in the case of a bacterial plasmid or a neomycin resistance gene for a mammalian vector. Vectors may be used in vitro, for example for the production of RNA or used to transfect or transform a host cell. The vector may also be adapted to be used in vivo, for example in a method of naked DNA vaccination or gene therapy. A further embodiment of the invention provides host cells transformed or transfected the vectors for the replication and expression of polynucleotides of the invention, including the DNA of GS, the open reading frames thereof and other corresponding ORFs particularly ORFs B, C, E and F from Mtb. The cells will be chosen to be compatible with the said vector and may for example be bacterial, yeast, insect or mammalian.

- 23 -

Expression vectors are widely available in the art and can be obtained commercially. Mammalian expression vectors may comprise a mammalian or viral promoter. Mammalian promoters include the metallothionien promoter. Viral promoters include promoters from adenovirus, the SV40 large T promoter and retroviral LTR promoters. Promoters compatible with insect cells include the polyhedrin promoter. Yeast promoters include the alcohol dehydrogenase promoter. Bacterial promoters include the β -galactosidase promoter.

The expression vectors may also comprise enhancers, and in the case of eukaryotic vectors polyadenylation signal sequence downstream of the coding sequence being expressed.

Polypeptides of the invention may be expressed in suitable host cells, for example bacterial, yeast, plant, insect and mammalian cells, and recovered using standard purification techniques including, for example affinity chromatography, HPLC or other chromatographic separation techniques.

Polynucleotides according to the invention may also be inserted into the vectors described above in an antisense orientation in order to provide for the production of antisense RNA. Antisense RNA or other antisense polynucleotides or ligands may also be produced by synthetic means. Such antisense polynucleotides may be used in a method of controlling the levels of the proteins encoded by the ORFs of the invention in a mycobacterial cell.

- Polynucleotides of the invention may also be carried by vectors suitable for gene therapy methods. Such gene therapy methods include those designed to provide vaccination against diseases caused by pathogenic mycobacteria or to boost the immune response of a human or animal infected with a pathogenic mycobacteria.
- 30 For example, Ziegner et al, AIDS, 1995, 9;43-50 describes the use of a replication defective recombinant amphotropic retrovirus to boost the immune response in patients with HIV infection. Such a retrovirus may be modified to carry a polynucleotide encoding a polypeptide or fragment thereof of the invention and the

15

20

25

35

retrovirus delivered to the cells of a human or animal subject in order to provide an immune response against said polypeptide. The retrovirus may be delivered directly to the patient or may be used to infecte cells ex-vivo, e.g. fibroblast cells, which are then introduced into the patient, optionally after being inactivated. The cells are desirably autologous or HLA-matched cells from the human or animal subject.

Gene therapy methods including methods for boosting an immune response to a particluar pathogen are disclosed generally in for example WO95/14091, the disclosure of which is incoporated herein by reference. Recombinant viral vectors include retroviral vectors, adenoviral vectors, adeno-associated viral vectors, vaccinia virus vectors, herpes virus vectors and alphavirus vectors. Alpha virus vectors are described in, for example, WO95/07994, the disclosure of which is incorporated herein by reference.

Where direct administration of the recombinant viral vector is contemplated, either in the form of naked nucleic acid or in the form of packaged particles carrying the nucleic acid this may be done by any suitable means, for example oral administration or intravenous injection. From 10⁵ to 10⁸ c.f.u of virus represents a typical dose, which may be repeated for example weekly over a period of a few months. Administration of autologous or HLA-matched cells infected with the virus may be more convenient in some cases. This will generally be achieved by administering doses, for example from 10⁵ to 10⁸ cells per dose which may be repeated as described above.

The recombinant viral vector may further comprise nucleic acid capable of expressing an accessory molecule of the immune system designed to increase the immune response. Such a molecule may be for example and interferon, particularly interferon gamma, an interleukin, for example IL-1 α , IL-1 β or IL-2, or an HLA class I or II molecule. This may be particularly desirable where the vector is intended for use in the treatment of humans or animals already infected with a mycobacteria and it is desired to boost the immune response.

30

E. Antibodies.

The invention also provides monoclonal or polyclonal antibodies to polypeptides of the invention or fragments thereof. The invention further provides a process for the production of monoclonal or polyclonal antibodies to polypeptides of the invention. Monoclonal antibodies may be prepared by conventional hybridoma technology using the polypeptides of the invention or peptide fragments thereof, as immunogens. Polyclonal antibodies may also be prepared by conventional means which comprise inoculating a host animal, for example a rat or a rabbit, with a polypeptide of the invention or peptide fragment thereof and recovering immune serum.

In order that such antibodies may be made, the invention also provides polypeptides of the invention or fragments thereof haptenised to another polypeptide for use as immunogens in animals or humans.

For the purposes of this invention, the term "antibody", unless specified to the contrary, includes fragments of whole antibodies which retain their binding activity for a polypeptide of the invention. Such fragments include Fv, F(ab') and $F(ab')_2$ fragments, as well as single chain antibodies. Furthermore, the antibodies and fragments thereof may be humanised antibodies, e.g. as described in EP-A-239400.

Antibodies may be used in methods of detecting polypeptides of the invention present in biological samples (where such samples include the human or animal body samples, and environmental samples, mentioned above) by a method which comprises:

- (a) providing an antibody of the invention;
- (b) incubating a biological sample with said antibody under conditions which allow for the formation of an antibody-antigen complex; and
- (c) determining whether antibody-antigen complex comprising said antibody is formed.

- 26 -

Antibodies of the invention may be bound to a solid support for example an immunoassay well, microparticle, dipstick or biosensor and/or packaged into kits in a suitable container along with suitable reagents, controls, instructions and the like.

5 Antibodies of the invention may be used in the detection, diagnosis and prognosis of diseases as descirbed above in relation to polypeptides of the invention.

F. Compositions.

The present invention also provides compositions comprising a polynucleotide or polypeptide of the invention together with a carrier or diluent. Compositions of the invention also include compositions comprising a nucleic acid, particularly and expression vector, of the invention. Compositions further include those carrying a recombinant virus of the invention.

15 Such compositions include pharmaceutical compositions in which case the carrier or diluent will be pharmaceutically acceptable.

Pharmaceutically acceptable carriers or diluents include those used in formulations suitable for inhalation as well as oral, parenteral (e.g. intramuscular or intravenous or transcutaneous) administration. The formulations may conveniently be presented in unit dosage form and may be prepared by any of the methods well known in the art of pharmacy. Such methods include the step of bringing into association the active ingredient with the carrier which constitutes one or more accessory ingredients. In general the formulations are prepared by uniformly and intimately bringing into association the active ingredient with liquid carriers or finely divided solid carriers or both, and then, if necessary, shaping the product.

For example, formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation isotonic with the blood of the intended recipient, and aqueous and non-aqueous sterile suspensions which may include suspending agents and thickening

cells such as M cells of the intestine after oral administration.

10

15

20

25

30

35

agents, and liposomes or other microparticulate systems which are designed to target the polynucleotide or the polypeptide of the invention to blood components or one or more organs, or to target

5 <u>G. Vaccines</u>.

In another aspect, the invention provides novel vaccines for the prevention and treatment of infections caused by Mptb, Mavs, other GS-containing pathogenic mycobacteria and Mtb in animals The term "vaccine" as used herein means an agent used to stimulate the immune system of a vertebrate, particularly a warm blooded vertebrate including humans, so as to provide protection against future harm by an organism to which the vaccine is directed or to assist in the eradication of an organism in the treatment of established infection. The immune system will be stimulated by the production of cellular immunity antibodies, desirably neutralizing antibodies, directed to epitopes found on or in a pathogenic mycobacterium which expresses any one of the ORFs of the invention. The antibody so produced may be any of the immunological classes, such as the immunoglobulins A, D, E, G or M. Vaccines which stimulate the production of IgA are interest since this is the principle immunoglobulin produced by the secretory system of warm-blooded animals, and the production of such antibodies will help prevent infection or colonization of the intestinal tract. IgM and IgG response will also be desirable for systemic infections such as Crohn's disease or tuberculosis.

Vaccines of the invention include polynucleotides of the invention or fragments thereof in suitable vectors and administered by injection of naked DNA using standard protocols. Polynucleotides of the invention or fragments thereof in suitable vectors for the expression of the polypeptides of the invention may be given by injection, inhalation or by mouth. Suitable vectors include M.bovis BCG, M.smegmatis or other mycobacteria, Corynebacteria, Salmonella or other agents according to established protocols.

20

25

35

Polypeptides of the invention or fragments thereof in substantially isolated form may be used as vaccines by injection, inhalation, oral administration or by transcutaneous application according to standard protocols. Adjuvants (such as Iscoms or polylactide-coglycolide encapsulation), cytokines such as IL-12 and other immunomodulators may be used for the selective enhancement of the cell mediated or humoral immunological responses. Vaccination with polynucleotides and/or polypeptides of the invention may be undertaken to increase the susceptibility of pathogenic mycobacteria to antimicrobial agents in vivo.

- 28 -

In instances wherein the polypeptide is correctly configured so as to provide the correct epitope, but is too small to be immunogenic, the polypeptide may be linked to a suitable carrier.

A number of techniques for obtaining such linkage are known in the art, including the formation of disulfide linkages using Nsuccinimidyl-3-(2-pyridylthio) propionate (SPDP) and succinimidyl 4-(N-maleimido-methyl)cyclohexane-1-carboxylate (SMCC) obtained from Pierce Company, Rockford, Illinois, (if the peptide lacks a sulfhydryl group, this can be provided by addition of a These reagents create a disulfide linkage cysteine residue). between themselves and peptide cysteine residues on one protein and an amide linkage through the epsilon-amino on a lysine, or other free amino group in the other. A variety of See, for example, disulfide/amide-forming agents are known. Immun Rev (1982) 62:185. Other bifunctional coupling agents form a thioether rather than a disulfide linkage. Many of these thioether-forming agents are commercially available and include reactive esters of 6-maleimidocaproic acid, 2-bromoacetic acid, 2-iodoacetic acid, 4-(N-maleimido-methyl)cyclohexane-1-carboxylic The carboxyl group can be activated by acid, and the like. combining them with succinimide or 1-hydroxyl-2-nitro-4-sulfonic Additional methods of coupling antigens acid, sodium salt. employs the rotavirus/"binding peptide" system described in EPO Pub. No. 259,149, the disclosure of which is incorporated herein by reference. The foregoing list is not meant to be exhaustive, and modifications of the named compounds can clearly be used.

Any carrier may be used which does not itself induce the production of antibodies harmful to the host. Suitable carriers are typically large, slowly metabolized macromolecules such as proteins; polysaccharides, such as latex functionalized Sepharose®, agarose, cellulose, cellulose beads and the like; polymeric amino acids, such as polyglutamic acid, polylysine, polylactide-coglycolide and the like; amino acid copolymers; and

are serum albumins, keyhole limpet hemocyanin, immunoglobulin molecules, thyroglobulin, ovalbumin, tetanus toxoid, and other proteins well known to those skilled in the art.

inactive virus particles. Especially useful protein substrates

The immunogenicity of the epitopes may also be enhanced by preparing them in mammalian or yeast systems fused with or assembled with particle-forming proteins such as, for example, that associated with hepatitis B surface antigen. See, e.g., US-A-4,722,840. Constructs wherein the epitope is linked directly to the particle-forming protein coding sequences produce hybrids which are immunogenic with respect to the epitope. In addition, all of the vectors prepared include epitopes specific to HBV, having various degrees of immunogenicity, such as, for example, the pre-S peptide.

In addition, portions of the particle-forming protein coding sequence may be replaced with codons encoding an epitope of the invention. In this replacement, regions which are not required to mediate the aggregation of the units to form immunogenic particles in yeast or mammals can be deleted, thus eliminating additional HBV antigenic sites from competition with the epitope of the invention.

Vaccines may be prepared from one or more immunogenic polypeptides of the invention. These polypeptides may be expressed in various host cells (e.g., bacteria, yeast, insect, or mammalian cells), or alternatively may be isolated from viral preparations or made synthetically.

In addition to the above, it is also possible to prepare live vaccines of attenuated microorganisms which express one or more

30

recombinant polypeptides of the invention. Suitable attenuated microorganisms are known in the art and include, for example, viruses (e.g., vaccinia virus), as well as bacteria.

The preparation of vaccines which contain an immunogenic polypeptide(s) as active ingredients, is known to one skilled in the art. Typically, such vaccines are prepared as injectables, or as suitably encapsulated oral preparations and either liquid solutions or suspensions; solid forms suitable for solution in, or suspension in, liquid prior to injestion or injection may also be prepared. The preparation may also be emulsified, or the protein encapsulated in liposomes. The active immunogenic often mixed with excipients which ingredients are pharmaceutically acceptable and compatible with the active ingredient. Suitable excipients are, for example, water, saline, dextrose, glycerol, ethanol, or the like and combinations thereof. In addition, if desired, the vaccine may contain minor amounts of auxiliary substances such as wetting or emulsifying agents, pH buffering agents, and/or adjuvants which enhance the effectiveness of the vaccine. Examples of adjuvants which may be effective include but are not limited to: aluminum hydroxide, N-acetyl-muramyl-L-threonyl-D-isoglutamine (thr-MDP), N-acetylnor-muramyl-L-alanyl-D-isoglutamine (CGP 11637, referred to as nor-MDP), N-acetylmuramyl-L-alanyl-D-isoglutaminyl-L-alanine-2-(1'-2'-dipalmitoyl-sn-glycero-3-hydroxyphosphoryloxy)-ethylamine (CGP 19835A, referred to as MTP-PE), and RIBI, which contains three components extracted from bacteria, monophosphoryl lipid A, trehalose dimycolate and cell wall skeleton (MPL+TDM+CWS) in a 2% squalene/Tween® 80 emulsion. The effectiveness of an adjuvant may be determined by measuring the amount of antibodies directed against an immunogenic polypeptide containing an antigenic sequence resulting from administration of this polypeptide in vaccines which are also comprised of the various adjuvants.

The vaccines are conventionally administered parenterally, by injection, for example, either subcutaneously or intramuscularly. Additional formulations which are suitable for other modes of administration include suppositories, oral formulations or as

enemas. For suppositories, traditional binders and carriers may include, for example, polyalkylene glycols or triglycerides; such suppositories may be formed from mixtures containing the active ingredient in the range of 0.5% to 10%, preferably 1% - 2%. Oral formulations include such normally employed excipients as, for example, pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate, and the like. These compositions take the form of solutions, suspensions, tablets, pills, capsules, sustained release formulations or powders and contain 10% - 95% of active ingredient, preferably 25% - 70%.

The proteins may be formulated into the vaccine as neutral or salt forms. Pharmaceutically acceptable salts include the acid addition salts (formed with free amino groups of the peptide) and which are formed with inorganic acids such as, for example, hydrochloric or phosphoric acids, or such organic acids such as acetic, oxalic, tartaric, maleic, and the like. Salts formed with the free carboxyl groups may also be derived from inorganic bases such as, for example, sodium, potassium, ammonium, calcium, or ferric hydroxides, and such organic bases as isopropylamine, trimethylamine, 2-ethylamino ethanol, histidine, procaine, and the like.

The vaccines are administered in a manner compatible with the dosage formulation, and in such amount as will be prophylactically and/or therapeutically effective. The quantity to be administered, which is generally in the range of $5\mu g$ to $250\mu g$, of antigen per dose, depends on the subject to be treated, capacity of the subject's immune system to synthesize antibodies, mode of administration and the degree of protection desired. Precise amounts of active ingredient required to be administered may depend on the judgement of the practitioner and may be peculiar to each subject.

The vaccine may be given in a single dose schedule, or preferably in a multiple dose schedule. A multiple dose schedule is one in which a primary course of vaccination may be with 1-10 separate doses, followed by other doses given at subsequent time intervals

25

required to maintain and or reenforce the immune response, for example, at 1-4 months for a second dose, and if needed, a subsequent dose(s) after several months. The dosage regimen will also, at least in part, be determined by the need of the individual and be dependent upon the judgement of the practitioner.

- 32 -

In a further aspect of the invention, there is provided an attenuated vaccine comprising a normally pathogenic mycobacteria which harbours an attenuating mutation in any one of the genes encoding a polypeptide of the invention. The gene is selected from the group of ORFs A, B, C, D, E, F, G and H, including the homologous ORFs B, C, E and F in Mtb.

The mycobacteria may be used in the form of killed bacteria or as a live attenuated vaccine. There are advantages to a live attenuated vaccine. The whole live organism is used, rather than dead cells or selected cell components which may exhibit modified or denatured antigens. Protein antigens in the outer membrane will maintain their tertiary and quaternary structures. Therefore the potential to elicit a good protective long term immunity should be higher.

The term "mutation" and the like refers to a genetic lesion in a gene which renders the gene non-functional. This may be at either the level of transcription or translation. The term thus envisages deletion of the entire gene or substantial portions thereof, and also point mutations in the coding sequence which result in truncated gene products unable to carry out the normal function of the gene.

A mutation introduced into a bacterium of the invention will generally be a non-reverting attenuating mutation. Non-reverting means that for practical purposes the probability of the mutated gene being restored to its normal function is small, for example less than 1 in 10^6 such as less than 1 in 10^9 or even less than 1 in 10^{12} .

20

25

30

An attenuated mycobacteria of the invention may be in isolated form. This is usually desirable when the bacterium is to be used for the purposes of vaccination. The term "isolated" means that the bacterium is in a form in which it can be cultured, processed or otherwise used in a form in which it can be readily identified and in which it is substantially uncontaminated by other bacterial strains, for example non-attenuated parent strains or unrelated bacterial strains. The term "isolated bacterium" thus encompasses cultures of a bacterial mutant of the invention, for example in the form of colonies on a solid medium or in the form of a liquid culture, as well as frozen or dried preparations of the strains.

In a preferred aspect, the attenuated mycobacterium further comprises at least one additional mutation. This may be a mutation in a gene responsible for the production of products essential to bacterial growth which are absent in a human or animal host. For example, mutations to the gene for aspartate semi-aldehyde dehydrogenase (asd) have been proposed for the production of attenuated strains of Salmonella. The asd gene is described further in Gene (1993) 129; 123-128. A lesion in the aspartate β -semialdehyde enzyme encoding the dehydrogenase would render the organism auxotrophic for the essential nutrient diaminopelic acid (DAP), which can be provided exogenously during bulk culture of the vaccine strain. this compound is an essential constituent of the cell wall for gram-negative and some gram-positive organisms and is absent from mammalian or other vertebrate tissues, mutants would undergo lysis after about three rounds of division in such tissues. Analogous mutations may be made to the attenuated mycobacteria of the invention.

In addition or in the alternative, the attenuated mycobacteria may carry a recA mutation. The recA mutation knocks out homologous recombination - the process which is exploited for the construction of the mutations. Once the recA mutation has been incorporated the strain will be unable to repair the constructed deletion mutations. Such a mutation will provide attenuated strains in which the possibility of homologous recombination to

30

35

with DNA from wild-type strains has been minimized. RecA genes have been widely studied in the art and their sequences are Further modifications may be made for additional available. safety.

The invention further provides a process for preparing a vaccine composition comprising an attenuated bacterium according to the invention process comprises (a) inoculating a culture vessel containing a nutrient medium suitable for growth of said bacterium; (b) culturing said bacterium; (c) recovering said 10 bacteria and (d) mixing said bacteria with a pharmaceutically acceptable diluent or carrier.

Attenuated bacterial strains according to the invention may be constructed using recombinant DNA methodology which is known per In general, bacterial genes may be mutated by a process of 15 targeted homologous recombination in which a DNA construct containing a mutated form of the gene is introduced into a host bacterium which it is desired to attenuate. The construct will recombine with the wild-type gene carried by the host and thus the mutated gene may be incorporated into the host genome to provide a bacterium of the present invention which may then be isolated.

The mutated gene may be obtained by introducing deletions into the gene, e.g by digesting with a restriction enzyme which cuts the coding sequence twice to excise a portion of the gene and then religating under conditions in which the excised portion is not reintroduced into the cut gene. Alternatively frame shift mutations may be introduced by cutting with a restriction enzyme which leaves overhanging 5' and 3' termini, filling in and/or trimming back the overhangs, and religating. Similar mutations may be made by site directed mutagenesis. These are only examples of the types of techniques which will readily be at the disposal of those of skill in the art.

Various assays are available to detect successful recombination. In the case of attenuations which mutate a target gene necessary for the production of an essential metabolite or catabolite

15

compound, selection may be carried out by screening for bacteria unable to grow in the absence of such a compound. Bacteria may also be screened with antibodies or nucleic acids of the invention to determine the absence of production of a mutated gene product of the invention or to confirm that the genetic lesion introduced - e.g. a deletion - has been incorporated into the genome of the attenuated strain.

The concentration of the attenuated strain in the vaccine will be formulated to allow convenient unit dosage forms to be prepared. Concentrations of from about 10⁴ to 10⁹ bacteria per ml will generally be suitable, e.g. from about 10⁵ to 10⁸ such as about 10⁶ per ml. Live attenuated organisms may be administered subcutaneously or intramuscularly at up to 10⁸ organisms in one or more doses, e.g from around 10⁵ to 10⁸, e.g about 10⁶ or 10⁷ organisms in a single dose.

The vaccines of the invention may be administered to recipients to treat established disease or in order to protect them against diseases caused by the corresponding wild type mycobacteria, such as inflammatory diseases such as Crohn's disease or sarcoidosis in humans or Johne's disease in animals. The vaccine may be administered by any suitable route. In general, subcutaneous or intramuscular injection is most convenient, but oral, intranasal and colorectal administration may also be used.

The following Examples illustrates aspects of the invention.

25 EXAMPLE 1

30

Tests for the presence of the GS identifier sequence were performed on $5\mu l$ bacterial DNA extracts (25 $\mu g/m l$ to 500 $\mu g/m l$) using polymerase chain reaction based on the oligonucleotide primers 5'-GATGCCGTGAGGAGGTAAAGCTGC-3' (Seq ID No. 40) and 5'-GATACGGCTCTTGAATCCTGCACG-3' (Seq ID No. 41) from within the identifier DNA sequences (Seq.ID Nos 1 and 2). PCR was performed for 40 cycles in the presence of 1.5 mM magnesium and an annealing temperature of 58°C. The presence or absence of the correct amplification product indicated the presence or absence

of GS identifier sequence in the corresponding bacterium. identifier sequence is shown to be present in all the laboratory and field strains of Mptb and Mavs tested. This includes Mptb isolates 0025 (bovine CVL Weybridge), 0021 (caprine, Moredun), (bovine, Moredun), 0139 (human, Chiodini 1984), 0209, 0208, 0211, 0210, 0212, 0207, 0204, 0206 (bovine, Whipple 1990). All Mptb strains were IS900 positive. The Mavs strains include 0010 and 0012 (woodpigeon, Thorel) 0018 (armadillo, Portaels) and 0034, 0037, 0038, 0040 (AIDS, Hoffner). All Mavs strains were One pathogenic M.avium strain 0033 (AIDS, IS902 positive. Hoffner) also contained GS identifier sequence. GS identifier sequence is absent from other mycobacteria including other M.malmoense, M.szulgai, M.gordonae, M. chelonei, M.avium, M. fortuitum, M. phlei, as well as E. coli, S. areus, Nocardia sp, Streptococcus sp. Shigella sp. Pseudomonas sp.

Example 2:

10

20

To obtain the full sequence of GS in Mavs and Mptb we generated a genomic library of Mavs using the restriction endonuclease EcoRI and cloning into the vector pUC18. This achieved a representative library which was screened with 32P-labelled identifier sequence yielding a positive clone containing a 17kbp insert. We constructed a restriction map of this insert and identified GS as fragments unique to Mavs and Mptb and not occurring in laboratory strains of M.avium. These fragments We identified GS were sub-cloned into pUC18 and pGEM4Z. The full nucleotide sequence contained within an 8kb region. was determined for GS on both DNA strands using primer walking and automated DNA sequencing. DNA sequence for GS in Mptb was obtained using overlapping PCR products generated using PwoDNA polymerase, a proofreading thermostable enzyme. The final DNA sequences were derived using the University of Wisconsin GCG gel assembly software package.

Example 3:

The DNA sequence of GS in Mavs and Mptb was found to be more than 99% homologous. The ORFs encoded in GS were identified using GeneRunner and DNAStar computer programmes. Eight ORFs were identified and designated GSA, GSB, GSC, GSD, GSE, GSF, GSG

Database comparisons were carried out against the and GSH. GenEMBL Database release version 48.0 (9/96), using the BLAST and BLIXEM programmes. GSA and GSB encoded proteins of 13.5kDa and 30.7kDa respectively, both of unknown functions. GSC encoded a protein of 38.4kDa with a 65% homology to the amino acid sequence of rfbD of V.cholerae, a 62% amino acid sequence homology to gmd of E.coli and a 58% homology to gca of Ps.aeruginosa which are all GDP-D-mannose dehydratases. Equivalent gene products in H.influenzae, S. dysenteriae, K.pneumoniae and Y.enterocolitica, N.gonorrhoea, 10 Salmonella enterica are all involved in 'O'-antigen processing known to be linked to pathogenicity. GSD encoded a protein of 37.1kDa which showed 58% homology at the DNA level to wcaG from a gene involved in the synthesis and regulation of capsular polysaccharides, also related to pathogenicity. was found to have a > 30% amino acid homology to rfbT of V. cholerae, involved in the transport of specific LPS components across the cell membrane. In V. cholerae the gene product causes a seroconversion from the Inaba to the Ogawa 'epidemic' strain. GSF encoded a protein of 30.2kDa which was homologous in the 20 range 25-40% at the amino acid level to several glucosyl transferases such as rfpA of K.pneumoniae, rfbB of K.pneumoniae, In E.coli an lgtD of H.influenzae, lsi of N.gonorrhoae. equivalent gene galE adds β -1-3 N-acetylglucosamine to galactose, the latter only found in 'O' and 'M' antigens which are also 25 related to pathogenicity. GSH comprising the ORFs GSH, and GSH2 encodes a protein totalling about 60kDa which is a putative transposase with a 40 - 43% homology at the amino acid level to the equivalent gene product of IS21 in E.coli. This family of insertion sequences is broadly distributed amongst gram negative bacteria and is responsible for mobility and transposition of genetic elements. An IS21- like element in B. fragilis is split either side of the β -lactamase gene controlling its activation and expression. We programmed an E.coli S30 cell-free extract with plasmid DNA containing the ORF GSH under the control of a the presence of a 35S-methionine, lac promoter in demonstrated the translation of an abundant 60kDa protein. The proteins homologous to GS encoded in other organisms are in general highly antigenic. Thus the proteins encoded by the ORFs

in GS may be used in immunoassays of antibody or cell mediated infections for diagnosing immuno-reactivity mycobacteria, particularly Mptb, Mavs and Mtb. Enhancement of host immune recognition of GS encoded proteins by vaccination using naked specific DNA or recombinant GS proteins, may be used in the prevention and treatment of infections caused by Mptb, Mavs and Mtb in humans and animals. Mutation or deletion of all or some of the ORFs A to H in GS may be used to generate attenuated strains of Mptb, Mavs or Mtb with lower pathogenicity for use as living or killed vaccines in humans and animals. Such vaccines are particularly relevant to Johne's disease in animals, to diseases caused by Mptb in humans such as Crohn's disease, and to the management of tuberculosis especially where the disease is caused by multiple drug-resistant organisms.





SEQUENCE LISTING

Seq. ID No.1

				**CCCCCCCC	AAACTATTCC	ACCTETECCE	GCTACGCAGT
						ACGTCTCCGC	
,						TGCAACGGTG	
5						GGCCGATGTT	
_						TCGCCTGGCC	
						CCTACGCATT	
	301	CGCTTTCGCG	CCACGTCGCA	GGCTTGCGGC	AGCATCCCGT	TCAGGTCTCC	TCATGGTCCG
	361	GTGTGGCACG	ACCACGCAAG	CTCGAACCGA	CTCGTTTCCC	AATTTCGCAT	GCTAATATCG
10	421	CTCGATGGAT	TTTTTGCGCA	ACGCCGGCTT	GATGGCTCGT	AACGTTAGCA	CCGAGATGCT
						GAAGCATACG	
	541	TGTTATTGAT	GTCGGTGCTA	ACTCCGGCCA	GTTCGGTAGC	GCTTTGCGTC	GTGCAGGATT
	601	CAAGAGCCGT	ATCGTTTCCT	TTGAACCTCT	TTCGGGGCCA	TTTGCGCAAC	TAACGCGCAA
	661	GTCGGCATCG	GATC -3'				

				_
15	Sea.	ID	No.	. 2

<u>narna tangon</u>

5'-	- 1	GATCCGATGC	CGACTTGCGC	GTTAGTTGCG	CAAATGGCCC	CGAAAGAGGT	TCAAAGGAAA
	61	CGATACGGCT	CTTGAATCCT	GCACGACGCA	AAGCGCTACC	GAACTGGCCG	GAGTTAGCAC
1	121	CGACATCAAT	AACAACGTTG	ACTCCGTATG	CTTCGACTTG	GTTTACTAAT	AGGCGCTTTC
-	181	GTTCGGAGTG	GCGCAGCATC	TCGGTGCTAA	CGTTACGAGC	CATCAAGCCG	GCGTTGCGCA
20 :	241	AAAAATCCAT	CGAGCGATAT	TAGCATGCGA	AATTGGGAAA	CGAGTCGGTT	CGAGCTTGCG
20	201	TEGTEGTECE	ACACCGGACC	ATGAGGAGAC	CTGAACGGGA	TGCTGCCGCA	AGCCTGCGAC
	161	CTGCCGCGAA	AGCGGTCAAC	TGATAATGCG	TAGGGGCGAG	CCAAGCCAGG	CTATGGGATA
	421	CTCACATACC	CCATGCGCCT	CGCCGGCCAG	GCGACTGCCG	TACCACTCGA	TGGCAGGTCG
•	421	CICACATAGG	CCGCCGGAG	GGATAACATC	GGCCGGCCCG	CAGCTTTACC	TCCTCACGGC
05	48T	-macaca mm	CTCCCTACC	ACGACACCGT	TGCATCGATG	CCCTCTTTCA	GTGCGATTCG
25	541	ATCGGCATII	GIGCGGIACC	aconcacco:	A COMOCA ATA	CTTTCCGCGG	GGTTCCATCG
!	601	CGGGCGCCAA	CCCAACTGCG	TAGCGCGGAG	ACGICCAMIA	9111900000	GGTTCCATCG
,	661	GGTTTAGTTG	GATC -3'				






	1			CGTCGAACTC		
	51			GCGGTGCCGA		
	101			GATGTATGAC		
5	151			CCGCCCCGCT '		
	201			GCGTGCTCGA		
	251			CTCACGGTTC		
	301	CGGGCTCGAA	AATCGGGACA	CTGCCTGCGA	GGAGCAAATC	GATCTTGGCC
	351	TGATCGATAT	CGACACAGAC	GACATCGTTG	CCGCTATCCG	CGAGACAGGC
10	401	GCCCGTGACG	AGGCCTACAT	AGCCTGATCC	GACCACCGAA	ATTTTCAAGA
	451	TGACCCCTTC	AAGTCCCCGA	TCGGTCGACG	ACCATACTGC	CGCAACTCTG
	501	TACCCTCCGT	GGGTAATTCG	CATGTCGCGT	TCGTAAGGAG	CAGCCAGCGA
	551	GTCGGGGACG	TTCGGTGAGA	GAGTCGCAGG	ACTACGAGGT	TGCCGGTGCG
	601	ATACATCACA	GTGTTGCGTC	TGTCGGCAAC	GATGCAGCAA	GAACCCACGG
15	651	GGCAGCCCTG	AACTGCGCGC	ATGACCGGTC	CTTGTCCTGG	CACCTTTGAT
	701	CGGCCACCGC	TTCCATGCGA	ACATGACCGG	AATCCATAGC	GCGTGGTCAA
	751	GCAGCGGGGA	GGTAGACGTC	GGTGTCATCT	GCTCCAACCG	TGTCGGTGAT
	801	AACGATTTCG	CTGAACGATC	TCGAGGGATT	GAAAAGCACC	GTGGAGAGCG
	851	TTCGCGCGCA	GCGCTATGGG	GGGCGAATCG	AGCACATCGT	CATCGACGGT
20	901			GGAGTATCTG		
	951			ACAACGGGAG		
	1001	GCATTGCCCA	TTCGTCGGGC	GACCTGTTGT	GGTTTATGCA	CTGCACGGAT
	1051			AGTCGCTTCC		
	1101			TGTGGGGTTA		
25	1151	GACTCGACGG	CAAACCACTT	TTCCCTCGGC	CGTACGGCTA	TATGCCGTTT
	1201			CGGCGCGACG		
	1251			CCAAGTTGGG		
	1301			TTCATCTACC		
	1351			GGTTTGCGAC		
30	1401			AGCACTATCG		
	1451			CTGGGTGGGC		
	1501			GATTCGGGCC		
	1551			AGTTCGCCAG		
	1601			GCCTGACCTG		
35	1651			CGGCCAGGTG		
•	1701			GGATCCCTTG		
	1751			GCTGGGCGG		
		AAGTGAGAAC				
		ATGGTTCCTA				
40		GGGCTCGTTC				
		CTACGTTGAC				
		ACCTCACTGA				
		GATGAGGTCT				
		CGAGCCAGTG				
45		TGGAAGCAGT				
••		TCGTCGGAGA				
						TCGTACTGGA
		CGACTCGCAA				
						TGACCCGAAA
50	2401					A TCGGAGGTCT
JU						GCCCGAATAT
						G ATGACTACGT
	2301	01CON0007				

5

10

15

50



CTGCAAGCCT GCCTCGGAAG CATCGTCGGG CAGACCTACC GGGAAGTGGA

5051 AGTGGTCCTT GTCGACGGCG GTTCGACCGA TCGGACCCTC GACATCGCGA 5101 ACAGTTTCCG CCCGGAACTC GGCTCGCGAC TGGTCGTTCA CAGCGGGCCC 5151 GATGATGGCC CCTACGACGC CATGAACCGC GGCGTCGGCG TGGCCACAGG

	5201	CGAATGGGTA	CTTTTTTTAG	GCGCCGACGA	CACCCTCTAC	GAACCAACCA
	5251	CGTTGGCCCA	GGTAGCCGCT	TTTCTCGGCG	ACCATGCGGC	AAGCCATCTT
	5301	GTCTATGGCG	ATGTTGTGAT	GCGTTCGACG	AAAAGCCGGC	ATGCCGGACC
	5351	TTTCGACCTC	GACCGCCTCC	TATTTGAGAC	GAATTTGTGC	CACCAATCGA
5	5401	TCTTTTACCG	CCGTGAGCTT	TTCGACGGCA	TCGGCCCTTA	CAACCTGCGC
	5451	TACCGAGTCT	GGGCGGACTG	GGACTTCAAT	ATTCGCTGCT	TCTCCAACCC
	5501	GGCGCTGATT	ACCCGCTACA	TGGACGTCGT	GATTTCCGAA	TACAACGACA
	5551	TGACCGGCTT	CAGCATGAGG	CAGGGGACTG	ATAAAGAGTT	CAGAAAACGG
	5601	CTGCCAATGT	ACTTCTGGGT	TGCAGGGTGG	GAGACTTGCA	GGCGCATGCT
10	5651	GGCGTTTTTG	AAAGACAAGG	AGAATCGCCG	TCTGGCCTTG	CGTACGCGGT
	5701	TGATAAGGGT	TAAGGCCGTC	TCCAAAGAAC	GAAGCGCAGA	ACCGTAGTCG
	5751	CGGATCCACA	TTGGACTTCT	TTAACGCGTT	TGCGTCCTGA	TCCACCTTTC
	5801	AAGCCCGTTC	CGCGTAACGC	GGCGCGCAGA	GAGTGGTCGC	ATATCGCATC
	5851	ACTGTTCTCG	TGCCAGTGCT	TGGAAAGCGT	CGAGCACTCT	GGTTCGCGTT
15	5901	CTTGACGTTC	GCGCCCGCTC	CTAGAGGTAG	CGTGTCACGT	GACTGAAGCC
	5951	AATGAGTGCA	ACTCGGCGTC	GCGAAAGGTT	TCAGTCGCGG	TTGAGCAAGA
	6001	CACCGCAAGA	CTACTGGAGT	GCGTGCACAA	GCGCCTCCAG	CTCGCGGCTG
	6051	AAAGCGGATG	CAAAGGGATT	CGAAGCTTGA	GCAACATGCG	AAGGGGAGAA
	6101	CGGCCTATGA	GGCTGGGACA	GGTTTTCGAT	CCGCGCGCGA	ATGCACTGTC
20	6151	AATGGCCAAG	TAGAAGTCCC	CGCTGGTGGC	CAGCAGAAGT	CCCCACTCCG
	6201	CTGCGGGTGG	TTGGCTAATT	CTTGGCGGCT	CCCTTCTTGT	GGTCGGCGTG
	6251	GCGCATCCGG	TAGGACTCGC	CGGAGGTGAC	GACGATGCTG	GCGTGGTGCA
	6301	GCAGCCGATC	GAGGATGCTG	GCGGCGGTGG	TGTGCTCGGG	CAGGAATCGC
	6351	CCCCATTGTT	CGAAGGGCCA	ATGCGAGGCG	ATGGCCAGGG	AGCGGCGCTC
25	6401	GTAGCCGGCA	GCCACGAGCC	GGAACAACAG	TTGAGTCCCG	GTGTCGTCGA
	6451	GCGGGGCGAA	GCCGATCTCG	TCCAAGATGA	CCAGATCCGC	GCGGAGCAGG
	6501	GTGTCGATGA	TCTTGCCGAC	GGTGTTGTCG	GCCAGGCCGC	GGTAGAGGAC
	6551	CTCGATCAGG	TCGGCGGCGG	TGAAGTAGCG	GACTTTGAAT	CCGGCGTGGA
	6601	CGGCAGCGTG	CCCGCAGCCG	ATGAGCAGGT	GACTTTTGCC	CGTACCAGGT
30	6651	GGGCCAATGA	CCGCCAGGTT	CTGTTGTGCC	CGAATCCATT	CCAGGCTCGA
	6701	CAGGTAGTCG	AACGTGGCTG	CGGTGATCGA	CGATCCGGTG	ACGTCGAACC
	6751	CGTCGAGGGT	CTTGGTGACC	GGGAAGGCTG	CGGCCTTGAG	ACGGTTGGCG
	6801	GTGTTGGAGG	CATCGCGGGC	AGCGATCTCG	GCCTCAACCA	ACGTCCGCAG
	6851			GTTGCGTCTT		
35	6901			GCCAGCTTCA		
	6951			CGCCGAGGAC		
						AGGCCTCCAA
				GCAGATCGAG		
-	7101			GCGCCGGCGG		
40	7151			CGCAACCGCC		
						TCGAGTTCGG
				ATCGCAGCAG		
						TTTTCGGGCG
	7351					TCATCGAGGA
45	7401					GATCACACCG
	7451			CAGGGCGCCA		
	7501					GAGCCGTAAC
	7551					CGAGCCGATC
	7601					CGTCAACCAA
50	7651					GCATTGACCT
	7701					GTCGACCTGC
	7751					GGTCGTCCTG
	7801	AGCGTAGCCA	CAGAGGTTC	r ccacgatgcc	CITCGATIG	GGATCCGCAC



7851 CGTGGCAGAA GTCCGGAACG AAGCCATAGT GGGACGCGAA TCGCACATAA
7901 TCCGGTGTTG GAACAACAAC ATTGGCGACG ACACCACCTT TGAGGCAGCC

- 43 -

7951 CATCCGGTCG GCCAGGATCT TGGCCGGAAC CCCACCGATC GCCTC

Seq. ID No.4

In the tent of the tent to

5	1	TTCTACTGCC	TGACCTGAGC	AGCGCCGAGG	CGCGCAGCGC	GATCACTGCG .	ACCTGAATGG
3	61	CCAGGTGGAA	AGCGCCACCG	ATCCCGGCAC	CGAGTGCCTG	ACGATTCGGA	TCCCTTGCAC
	121	CACAACGAGA	GTGAGACCGC	CATGATGACG	AAATATCGGC	TGGGCGGAGT	CAACGCCGGA
	121	GTGACAAAAG	TGAGAACCCG	GTGAAGCGAG	CGCTTATAAC	AGGGATCACG	GGGCAGGATG
	241	GTTCCTACCT	CGCCGAGCTA	CTACTGAGCA	AGGGATACGA	GGTTCACGGG	CTCGTTCGTC
10	241	GAGCTTCGAC	GTTTAACACG	TCGCGGATCG	ATCACCTCTA	CGTTGACCCA	CACCAACCGG
10	361	GCGCGCGCTT	CTTCTTGCAC	TATGCAGACC	TCACTGACGG	CACCCGGTTG	GTGACCCTGC
	421	TCAGCAGTAT	CGACCCGGAT	GAGGTCTACA	ACCTCGCAGC	GCAGTCCCAT	GTGCGCGTCA
	421	GCTTTGACGA	GCCAGTGCAT	ACCGGAGACA	CCACCGGCAT	GGGATCGATC	CGACTTCTGG
	401	AAGCAGTCCG	CCTTTCTCGG	GTGGACTGCC	GGTTCTATCA	GGCTTCCTCG	TCGGAGATGT
45	541	TCGGCGCATC	TCCCCCACCG	CAGAACGAAT	CGACGCCGTT	CTATCCCCGT	TCGCCATACG
15	601	GCGCGGCCAA	CCTCTTCTCG	TACTGGACGA	CTCGCAACTA	TCGAGAGGCG	TACGGATTAT
	991	TCGCAGTGAA	TCCCATCTTG	TTCAACCATG	AGTCCCCCG	GCGCGGCGAG	ACTTTCGTGA
	721	CCCGAAAGAT	CACCCCCCCCC	GTGGCGCGCA	TCCGAGCTGG	CGTCCAATCG	GAGGTCTATA
	781	TGGGCAACCT	CALGCGIGCC	CCCACTGGG	GCTACGCGCC	CGAATATGTC	GAGGGGATGT
	841	GGAGGATGTT	CGAIGCGAIC	CAACCTGATG	ACTACGTCCT	GGCGACAGGG	CGTGGTTACA
20	901	CCGTACGTGA	GCAAGCGCC1	CONCUMENT	ACCACGTCGG	GCTCGACTGG	CAAAAGCACG
	961	TCAAGTTTGA	GTTCGCTCAA	TTECECCCCA	CCGAGGTCGA	TTCGCTAGTA	GGAGATGCCG
	1021	TCAAGTTIGA ACAGGGCGGC	CGACCGCIAI	recreation.	CTTCGGTTCA	TACTGGTGAA	CTCGCGCGCA
	1081	ACAGGGCGGC TCATGGTGGA	CCAGTCACTC	. GGCIGGAAAG	ACTCCGATGG	CACACCATGG	ATCGACACGC
	1141	TCATGGTGGA CGATGTTGCC	CGCGGACATC	. GCCGCGICGG	GACGACTACA	CCTGGGCCTC	TGGACCGCGC
25	1201	CGATGTTGCC AACGCCCGTG	TGGTTGGGG	, GGAGIAAGII	COTESTOS	TCAGCGCTCG	TACGTAGATT
	1261	. AACGCCCGTG . TGAGGCCGAG	TATATUGUCU	, GICAICGGGG	CCCATCACGC	CATGAGATTG	ATCTGACGGA
	1321	. TGAGGCCGAG . CCGAGCCGCA	GGGTTCACCA	ATCTCATIGI	CACAAGACCA	CAGGTGATCA	TCGATGCGGC
	1381	. CCGAGCCGCA . CGCACGGGTC	ACGTTTGATT	TIGIGICIGA	CACARGACCA	GCGGACTTCT	TGTCCGAAAA
	1441	CGCACGGGTC CCTCCGAATC	GGCGGCATCA	TGGCGAATAF	. ACCTETCEC	CTGCGTGTGC	CGCGGCTCCT
30	1501	L CCTCCGAATO	CAGACCAAT	r TGCTCGACGC	CTACCCTCC	CAACCTATCC	ACGAGAGTGC
	1561	TTTCCTCGGT	TCGTCATGC	TCTACCCGA	CONCECTAN	GCGATCGCCA	AGATCGCCGG
	1621	TTTATTGACT	GGCCCTTGC	AGCCCACCA	A CGACGCGIA	CCTGGATCT	CTGCGATGCC
	168:	TATCCTGCA	GTTCAGGCG	TTAGGCGCC	RECECCETO	GCCTCGCATC	TCTTGCCGGC
	174:	GACTAACCT	TACGGACCC	G GCGACAACI.	r macracarco	CANGAGGTGA	CGAATTGGGG
35	180	L GCTCATCCG1	CGATATGAG	G AAGCCAAAG	TGGIGGIGG	r creeceaece	CGAATTGGGG
	186	l GACCGGTAC	. CCGCGGCGC	G AACTTCIGC	A IGICGACGA.	COCACCGGGG	CATGCCTGTT
	192	1 CCTTTTGGA	CATTTCGAT	G GTCCGAACC	A CGTCAACGT	TACATCGGCG	TCGATCACAG
	198	1 CATTAGCGA	ATCGCAGAC	A TGGTCGCTA	C GGCGGTGGG	C CACGTCTCCG	AAACACGTTG
	204	1 GGATCCAAC	r AAACCCGAT	G GAACCCCGC	G CAAACIAII	G CATGCAACGG	CGCTACGCGA
40	210	1 GTTGGGTTG	G CGCCCGCGA	A TCGCACTGA	A AGACGGCAI	C CCCCCCATC	TTATCCTCC
	216	1 CCGCACAAA	T GCCGATGCC	G TGAGGAGGT	A AAGCIGCGG	C ACTCCCCTCC	TTATCCCTCC
	222	1 GGCCGGACG	G GTAGGGCGA	C CTGCCATCG	A GIGGIACGG	C CCCCTACGC	CCGGCGAGGC
	228	1 GCATGGCCT	A TGGGAGTAT	C CCATAGCCT	G GCTTGGCTC	c contractor	TTATCAGTTG
	234	1 ACCGCTTTC	G CGCCAGCTC	G CAGGCTCGC	G GCAGCATCC	C GIICAGGIC:	CCTCATGGTC
45	240	1 CGGTGTGGC	A CGACCACGO	A AGCTCGAAC	C GACTCGTTT	C CCAATITCG	ATGCTAATAT
	246	1 CGCTCGATG	G ATTITITGO	G CAACGCCGG	C TTGATGGC:	C GIAACGIIA	G CACCGAGATG
	252	1 CTGCGCCAC	T TCGAACGAA	LA GCGCCTATI	A GTAAACCAA	TCAAAGCAT	A CGGAGTCAAC
	258	1 GTTGTTATI	G ATGTCGGTC	SC TAACTCCGC	C CAGTICGGI	A GUGUTTIGU	G TCGTGCAGGA
	264	1 TTCAAGAGC	C GTATCGTT	C CTTTGAACO	T CTTTCGGGC	C CHILIGCGC	A ACTAACGCGC
50	270	1 GAGTCGGCA	T CGGATCCA	CT ATGGGAGTO	T CACCAGTAT	G CCCIMOGCO	A CGCCGATGAG



- 44 -

				40			
	2761	ACGATTACCA	TCAATGTGGC	AGGCAATGCG	GGGGCAAGTA	GTTCCGTGCT	GCCGATGCTT
			AAGATGCCTT				
	2881	CACCGCCTTG	ATTCGGTTGC	ATCAGAATTT	CTGAACCCTA	CCGATGTTAC	TTTCCTGAAG
			AGGGTTTCGA				
5	3001	AGCTGCGTCG	GCATGCAACT	CGAACTTTCT	TTTATTCCGT	TGTACGAAGG	TGACATGCTG
	3061	ATTCATGAAG	CGCTTGAACT	TGTCTATTCC	CTAGGTTTCA	GACTGACGGG	TTTGTTGCCC
	3121	GGATTTACGG	ATCCGCGCAA	TGGTCGAATG	CTTCAAGCTG	ACGGCATTTT	CTTCCGTGGG
	3181	GACGATTGAC	ATAAATGCTT	GCGTCGGCAC	CCTGCCGGTA	TCCAAACGGG	CGATCTGGTG
	3241	AGCCGGCCTC	CCGGGCACCT	AATCGACTAT	CTAAATTGAG	GCGGCCGCGA	CGTGCGGCAC
10	3301	GAACAGGTGG	CCGGCTGCTA	GCGTTACACA	CGTCATGACT	GCGCCAGTGT	TCTCGATAAT
	3361	TATCCCTACC	TTCAATGCAG	CGGTGACGCT	GCAAGCCTGC	CTCGGAAGCA	TCGTCGGGCA
	3421	GACCTACCGG	GAAGTGGAAG	TGGTCCTTGT	CGACGGCGGT	TCGACCGATC	GGACCCTCGA
			AGTTTCCGCC				
			TACGACGCCA				
15	3601	TTTTTTAGGC	GCCGACGACA	CCCTCTACGA	ACCAACCACG	TTGGCCCAGG	TAGCCGCTTT
			CATGCGGCAA				
							ATTTGTGCCA
	3781	CCAATCGATC	TTTTACCGCC	GTGAGCTTTT	CGACGGCATC	GGCCCTTACA	ACCTGCGCTA
	3841	CCGAGTCTGG	GCGGACTGGG	ACTTCAATAT	TCGCTGCTTC	TCCAACCCGG	CGCTGATTAC
20							GCATGAGGCA
	3961	GGGGACTGAT	AAAGAGTTCA	GAAAACGGCT	GCCAATGTAC	TTCTGGGTTG	CAGGGTGGGA
	4021	GACTTGCAGG	CGCATGCTGG	CGTTTTTGAA	AGACAAGGAG	AATCGCCGTC	TGGCCTTGCG
	4081	TACGCGGTTG	ATAAGGGTTA	AGGCCGTCTC	CAAAGAACGA	AGCGCAGAAC	CGTAGTCGCG
	4141	GATCCACATT	GGACTTCTTT	AACGCGTTTG	CGTCCTGATC	CACCITICAA	CCCCGTTCCG
25	4201	CGTGACGCGG	CGCGCAGAGA	GTGGTCGCAT	ATCGCGTCAC	TGTTCTCGTG	CCAGTGCTTG
	4261	GAAAGCGTCG	AGCACTCTGG	TTCGCGTTCT	TGACGTTCGC	GCCCGCCCCT	AGAGGTAGCG
	4321	TGTCACGTGA	CTGAAGCCAA	TGAGTGCAAC	TOGGOGTOGO	GAAAGGTTTC	AGTCGCGGTT
	4381	. GAGCAAGACA	CCGCAAGACT	ACTGGAGTGC	GTGCACAAGC	GCCTCCAGCT	CACGG
		_					

Seq. ID No.5

1 atgategetg tgatetggte ggeggtgeeg acaggaaceg tegaettgte gaegateace
61 ttgtaceggt egatgtatga eccaatgteg teegcaaceg agaagaegta egteaggtee
121 geegeeege ttteacecat gggegteggg acggegatga aaatgaegte egegtgeteg
181 atteegegtt geeggteggt ggtgaagtea ateageeegt teteaceggtt eetegeaate
241 aacteecaac eegggetega aaategggae actgeetgeg aggageaaat egatettgge
35 301 etgategata tegacacaga egacategtt geegetatee gegagacagg egecegtgae
361 gaggeetaca tageetga

Seq. ID No.6

1 MIAVIWS AV PTGTVDLSTITLYRS MYDPMS
31 SATEKTYVRS AAPLS PMGVGTAMKMTS ACS
61 IPRCRS VVKS IS PFSRFLAL NS QPGLENRD
91 TACEEQIDLGLIDIDTDDIVAAIRETGARD
121 EAYIA



1 gtgtcatctg ctccaaccgt gtcggtgata acgatttcgc tgaacgatct cgagggattg 61 aaaagcaccg tggagagcgt tegegegeag egetatgggg ggegaatega geacategte 121 atcgacggtg gatcgggcga cgccgtcgtg gagtatctgt ccggcgatcc tggctttgca 181 tattggcaat ctcagcccga caacgggaga tatgacgcga tgaatcaggg cattgcccat 5 241 togtogggeg acctgttgtg gtttatgcac tocacggate gttteteega tecagatgca 301 gtcgcttccg tggtggaggc gctctcgggg catggaccag tacgtgattt gtggggttac 361 gggaaaaaca accttgtcgg actcgacggc aaaccacttt tccctcggcc gtacggctat 421 atgccgttta agatgcggaa atttctgctc ggcgcgacgg ttgcgcatca ggcgacattc 481 ttcggcgcgt cgctggtagc caagttgggc ggttacgatc ttgattttgg actcgaggcg 10 541 gaccagetgt teatetaceg tgccgcaeta atacggeete cegteacgat cgaccgcgtg 601 gtttgcgact tcgatgtcac gggacctggt tcaacccagc ccatccgtga gcactatcgg 661 accetgegge ggetetggga cetgeatgge gactaceege tgggtgggeg cagagtgteg 721 tgggcttact tgcgtgtgaa ggagtacttg attcgggccg acctggccgc attcaacgcg 781 gtaaagttct tgcgagcgaa gttcgccaga gcttcgcgga agcaaaattc atag 15

Seq. ID No.8

1 V S S A P T V S V I T I S L N D L E G L K S T V E S V R A Q

31 R Y G G R I E H I V I D G G S G D A V V E Y L S G D P G F A

61 Y W Q S Q P D N G R Y D A M N Q G I A H S S G D L L W F M H

20 91 S T D R F S D P D A V A S V V E A L S G H G P V R D L W G Y

121 G K N N L V G L D G K P L F P R P Y G Y M P F K M R K F L L

151 G A T V A H Q A T F F G A S L V A K L G G Y D L D F G L E A

181 D Q L F I Y R A A L I R P P V T I D R V V C D F D V T G P G

211 S T Q P I R E H Y R T L R R L W D L H G D Y P L G G R R V S

25 241 W A Y L R V K E Y L I R A D L A A F N A V K F L R A K F A R

271 A S R K Q N S

Seq. ID No.9

1 gtgaagcgag cgcttataac agggatcacg gggcaggatg gttcctacct cgccgagcta 61 ctactgagca agggatacga ggttcacggg ctcgttcgtc gagcttcgac gtttaacacg 121 tegeggateg atcaceteta egttgaceca caccaacegg gegegegett gttettgcac 30 181 tatgcagacc tcactgacgg cacccggttg gtgaccctgc tcagcagtat cgacccggat 241 gaggtetaca acetegeage geagteceat gtgegegtea getttgaega geeagtgeat 301 accggagaca ccaccggcat gggatcgatc cgacttetgg aagcagteeg cetttetegg 361 gtggactgcc ggttctatca ggcttcctcg tcggagatgt tcggcgcatc tccgccaccg 421 cagaacgaat cgacgccgtt ctatccccgt tegccatacg gegeggccaa ggtcttctcg 35 481 tactggacga ctcgcaacta tcgagaggcg tacggattat tcgcagtgaa tggcatcttg 541 ttcaaccatg agtccccccg gcgcggcgag actttcgtga cccgaaagat cacgcgtgcc 601 gtggcgcgca tccgagctgg cgtccaatcg gaggtctata tgggcaacct cgatgcgatc 661 cgcgactggg gctacgcgcc cgaatatgtc gaggggatgt ggaggatgtt gcaagcgcct 721 gaacctgatg actacgtcct ggcgacaggg cgtggttaca ccgtacgtga gttcgctcaa 40 781 gctgcttttg accatgtcgg gctcgactgg caaaagcgcg tcaagtttga cgaccgctat 841 ttgcgtccca ccgaggtcga ttcgctagta ggagatgccg acaaggcggc ccagtcactc 901 ggctggaaag cttcggttca tactggtgaa ctcgcgcgca tcatggtgga cgcggacatc 961 geogegtigg agigegatgg cacaccatgg ategacacge egatgitgee iggitiggge 45 1021 agagtaagtt ga



Seq. ID No.11

1 gtgaagegag egettataae agggateaeg gggeaggatg gtteetaeet egeegageta 15 61 ctactgagca agggatacga ggttcacggg ctcgttcgtc gagcttcgac gtttaacacg 121 tegeggateg atcaceteta egitgaecea caccaacegg gegegegett gitetigeae 181' tatgcagace teactgacgg cacceggttg gtgaccetge teagcagtat egacceggat 241 gaggtetaca acetegeage geagteceat gtgegegtea getttgaega geeagtgeat 301 accggagaca ccaccggcat gggatcgatc cgacttctgg aagcagtccg cctttctcgg 20 361 gtggactgcc ggttctatca ggcttcctcg tcggagatgt tcggcgcatc tccgccaccg 421 cagaacgaat cgacgccgtt ctatccccgt tcgccatacg gcgcggccaa ggtcttctcg 481 tactggacga ctcgcaacta tcgagaggcg tacggattat tcgcagtgaa tggcatcttg 541 ttcaaccatg agtccccccg gcgcggcgag actttcgtga cccgaaagat cacgcgtgcc 601 gtggegegea teegagetgg egteeaateg gaggtetata tgggeaacet egatgegate 25 661 cgcgactggg gctacgcgcc cgaatatgtc gaggggatgt ggaggatgtt gcaagcgcct 721 gaacetgatg actacgteet ggcgacaggg egtggttaca eegtacgtga gttegetcaa 781 getgettttg accaegtegg getegaetgg caaaageaeg teaagtttga egaeegetat 841 ttgcgcccca ccgaggtcga ttcgctagta ggagatgccg acagggcggc ccagtcactc 901 ggctggaaag cttcggttca tactggtgaa ctcgcgcgca tcatggtgga cgcggacatc 30 961 gccgcgtcgg agtgcgatgg cacaccatgg atcgacacgc cgatgttgcc tggttggggc 1021 ggagtaagtt ga

Seq. ID No.12



1 grgcgarggc acaccargga rcgacacgcc gargrrgccr ggrrggggca gagraagtrg 61 acgactacac ctgggcctct ggaccgcgca acgcccgtgt atatcgccgg tcatcggggg 121 ctggtcggct cagcgctcgt acgtagattt gaggccgagg ggttcaccaa tctcattgtg 181 cgatcacgcg atgagattga tctgacggac cgagccgcaa cgtttgattt tgtgtctgag 5 241 acaagaccac aggtgatcat cgatgcggcc gcacgggtcg gcggcatcat ggcgaataac 301 acctateceg eggaettett greegaaaac eteegaatee agaecaattt getegaegea 361 gctgtcgccg tgcgtgtgcc gcggctcctt ttcctcggtt cgtcatgcat ctacccgaag 421 tacgeteege aacetateea egagagtget ttattgactg gecetttgga geceaecaae 481 gacgcgtatg cgatcgccaa gatcgccggt atcctgcaag ttcaggcggt taggcgccaa 10 541 tatgggetgg egtggatete tgegatgeeg actaacetet aeggaceegg egacaaette 601 teccegteeg ggtegeatet ettgeeggeg etcateegte gatatgagga agecaaaget 661 ggtggtgcag aagaggtgac gaattggggg accggtactc cgcggcgcga acttctgcat 721 gtcgacgatc tggcgagcgc atgcctgttc cttttggaac atttcgatgg tccgaaccac 781 gtcaacgtgg gcaccggcgt cgatcacagc attagcgaga tcgcagacat ggtcgctaca 15 841 geggtggget acateggega aacaegttgg gatecaacta aaceegatgg aaceeegege 901 aaactattgg acgtctccgc gctacgcgag ttgggttggc gcccgcgaat cgcactgaaa 961 gacggcatcg atgcaacggt gtcgtggtac cgcacaaatg ccgatgccgt gaggaggtaa

Seq. ID No.14



1 grgcgarggc acaccargga rcgacacgcc gargrigccr ggrrggggcg gagraagtrg 61 acgactacac ctgggcctct ggaccgcgca acgcccgtgt atatcgccgg tcatcggggg 121 ctggtcggct cagcgctcgt acgtagattt gaggccgagg ggttcaccaa tctcattgtg 181 cgatcacgcg atgagattga tctgacggac cgagccgcaa cgtttgattt tgtgtctgag 5 241 acaagaccac aggtgatcat cgatgcggcc gcacgggtcg gcggcatcat ggcgaataac 301 acctatocog oggacttott gtoogaaaac otoogaatoo agaccaattt gotogacgca 361 getgtegeeg tgegtgtgee geggeteett tteeteggtt egteatgeat etaecegaag 421 tacgeteege aacetateea egagagtget ttattgactg gecetttgga geceaecaae 481 gacgcgtatg cgatcgccaa gatcgccggt atcctgcaag ttcaggcggt taggcgccaa 10 541 tatgggctgg cgtggatctc tgcgatgccg actaacctct acggacccgg cgacaacttc 601 teccegteeg ggtegeatet ettgeeggeg eteateegte gatatgagga agecaaaget 661 ggtggtgcag aagaggtgac gaattggggg accggtactc cgcggcgcga acttctgcat 721 gtcgacgatc tggcgagcgc atgcctgttc cttttggaac atttcgatgg tccgaaccac 781 gtcaacgtgg gcaccggcgt cgatcacagc attagcgaga tcgcagacat ggtcgctacg 15 841 gcggtgggct acatcggcga aacacgttgg gatccaacta aacccgatgg aaccccgcgc 901 aaactattgg acgtctccgc gctacgcgag ttgggttggc gcccgcgaat cgcactgaaa 961 gacggcatcg atgcaacggt gtcgtggtac cgcacaaatg ccgatgecgt gaggaggtaa

Seq. ID No.16

Seq. ID No.17

1 atggatttt tgcgcaacge cggcttgatg gctcgtaacg ttagtaccga gatgctgcgc
35 61 cacttcgaac gaaagcgcct attagtaaac caattcaaag catacggagt caacgttgtt
121 attgatgtcg gtgctaactc cggccagttc ggtagcgctt tgcgtcgtgc aggattcaacg
181 agccgtatcg tttcctttga acctctttcg gggccatttg cgcaactaac gcgcaagtcg
241 gcatcggatc cactatggga gtgtcaccag tatgccctag gcgacgccga tgagacgatt
301 accatcaatg tggcaggcaa tggcggggca agtagtccg tgctgccgat gcttaaaagt
40 361 catcaagatg cctttcctcc cgcgaattat attggcaccg aagacgttg actacacgg
421 cttgattcgg ttgcatcaga atttctgaac cctaccgatg ttactttcct gaagatcgac
481 gtacagggtt tcgagaagca ggttatcac ggcagtaagt caacgcttaa cgaaagctgc
541 gtcggcatgc aacttgtcaa ttccttatt ccgttgacg aaggtgacat gctgattcat
601 gaagcgcttg aacttgtcta ttccctaggt ttcagactga cgggtttgtt gcccggcttt
45 661 acggatccgc gcaatggtcg aatgctcaa gctgacggca ttttcttccg tggggacgat





1 M D F L R N A G L M A R N V S T E M L R H F E R K R L L V N
31 Q F K A Y G V N V V I D V G A N S G Q F G S A L R R A G F K
61 S R I V S F E P L S G P F A Q L T R K S A S D P L W E C H Q
91 Y A L G D A D E T I T I N V A G N A G A S S S V L P M L K S
121 H Q D A F P P A N Y I G T E D V A I H R L D S V A S E F L N
151 P T D V T F L K I D V Q G F E K Q V I T G S K S T L N E S C
181 V G M Q L E L S F I P L Y E G D M L I H E A L E L V Y S L G
211 F R L T G L L P G F T D P R N G R M L Q A D G I F F R G D D

10 Seq. ID No.19

5

15

20

1 atggatttt tgegeaaege eggettgatg getegtaaeg tageaeega gatgetgege 61 cacttegaae gaaagegeet attagtaaae caatteaaag cataeggagt caaegttgtt 121 attgatgteg gtgetaaete eggeeagtte ggtagegett tgegtegtge aggatteaag 181 ageegtateg ttteetttga acetettteg gggeeatttg egeaaetaae gegegagteg 241 geateggate cactatggga gtgteaeeag tatgeeetag gegaegeega tgagaegatt 301 aceateaatg tggeaggeaa tgegggggea agtagtteeg tgetgeegat gettaaaagt 361 cateaagatg eettteetee egegaattat attggeaeeg aagaegttge aataeaeege 421 ettgattegg ttgeateaga atteetgaae eetaeegatg ttaettteet gaagategae 481 gtaeagggtt tegagaagea ggttategeg ggeagtaagt eaaegettaa egaaagetge 541 gteggeatge aactegaeet tteetttatt eegttgaeg aaggtgaeat geeggatte 661 aeggateege geaatggteg aatgetteaa getgaeeggea ttteetteeg tggggaegat 721 tga

Seq. ID No.20

25

1- M D F L R N A G L M A R N V S T E M L R H F E R K R L L V N
31 Q F K A Y G V N V V I D V G A N S G Q F G S A L R R A G F K
61 S R I V S F E P L S G P F A Q L T R E S A S D P L W E C H Q
91 Y A L G D A D E T I T I N V A G N A G A S S S V L P M L K S
121 H Q D A F P P A N Y I G T E D V A I H R L D S V A S E F L N
151 P T D V T F L K I D V Q G F E K Q V I A G S K S T L N E S C
181 V G M Q L E L S F I P L Y E G D M L I H E A L E L V Y S L G
211 F R L T G L L P G F T D P R N G R M L Q A D G I F F R G D D





1 atgactgcgc cagtgttctc gataattatc cctaccttca atgcagcggt gacgctgcaa 61 gcctgcctcg gaagcatcgt cgggcagacc taccgggaag tggaagtggt ccttgtcgac 121 ggeggttega cegateggae cetegacate gegaacagtt teegeeegga acteggeteg 181 cgactggtcg ttcacagegg geoegatgat ggeecetaeg aegeeatgaa eegeggegte 5 241 ggcgtggcca caggcgaatg ggtacttttt ttaggcgccg acgacaccct ctacgaacca 301 accarging cocaggiage egethtetic ggegaccatg eggeaageca tethgictat 361 ggcgatgttg tgatgcgttc gacgaaaagc cggcatgccg gacctttcga cctcgaccgc 421 ctcctatttg agacgaattt gtgccaccaa tcgatctttt accgccgtga gcttttcgac 481 ggcatcggcc cttacaacct gcgctaccga gtctgggcgg actgggactt caatattcgc 10 541 tgcttctcca acceggeget gattaccege tacatggacg tegtgattte egaatacaac 601 gacatgaccg getteageat gaggeagggg actgataaag agtteagaaa acggetgeea 661 atgtacttct gggttgcagg gtgggagact tgcaggcgca tgctggcgtt tttgaaagac 721 aaggagaatc gccgtctggc cttgcgtacg cggttgataa gggttaaggc cgtctccaaa 15 781 gaacgaagcg cagaaccgta g

Seq. ID No.22

1 M T A P V F S I I I P T F N A A V T L Q A C L G S I V G Q T

31 Y R E V E V V L V D G G S T D R T L D I A N S F R P E L G S

61 R L V V H S G P D D G P Y D A M N R G V G V A T G E W V L F

91 L G A D D T L Y E P T T L A Q V A A F L G D H A A S H L V Y

121 G D V V M R S T K S R H A G P F D L D R L L F E T N L C H Q

151 S I F Y R R E L F D G I G P Y N L R Y R V W A D W D F N I R

181 C F S N P A L I T R Y M D V V I S E Y N D M T G F S M R Q G

211 T D K E F R K R L P M Y F W V A G W E T C R R M L A F L K D

25 241 K E N R R L A L R T R L I R V K A V S K E R S A E P

Seq. ID No.23

1 atgactgcgc cagtgttctc gataattatc cctaccttca atgcagcggt gacgctgcaa 61 gcctgcctcg gaagcatcgt cgggcagacc taccgggaag tggaagtggt ccttgtcgac 121 ggcggttcga ccgatcggac cctcgacatc gcgaacagtt tccgcccgga actcggctcg 181 cgactggtcg ttcacagcgg gcccgatgat ggcccctacg acgccatgaa ccgcggcgtc 30 241 ggcgtagcca caggcgaatg ggtacttttt ttaggcgccg acgacaccct ctacgaacca 301 accacgttgg cccaggtagc cgcttttctc ggcgaccatg cggcaagcca tcttgtctat 361 ggcgatgttg tgatgcgttc gacgaaaagc cggcatgccg gacctttcga cctcgaccgc 421 ctcctatttg agacgaattt gtgccaccaa tcgatctttt accgccgtga gcttttcgac 481 ggcatcggcc cttacaacct gcgctaccga gtctgggcgg actgggactt caatattcgc 35 541 tgcttctcca accoggogot gattaccogo tacatggacg togtgattto ogaatacaac 601 gacatgaccg getteageat gaggeagggg actgataaag agtteagaaa acggetgeea 661 atgtacttct gggttgcagg gtgggagact tgcaggcgca tgctggcgtt tttgaaagac 721 aaggagaatc geegtetgge ettgegtaeg eggttgataa gggttaagge egteteeaaa 40 781 gaacgaagcg cagaaccgta g

15

20

25





Seq. ID No.24

1 MTAPVFSIIIPTFNAAVTLQACLGSIVGQT Y R E V E V V L V D G G S T D R T L D I A N S F R P E L G S R L V V H S G P D D G P Y D A M N R G V G V A T G E W V L F LGADDTLYEPTTLAQVAAFLGDHAASHLVY 5 G D V V M R S T K S R H A G P F D L D R L L F E T N L C H Q SIFYRRELFDGIGPYNLRYRVWADWDFNIR C F S N P A L I T R Y M D V V I S E Y N D M T G F S M R Q G 211 TDKEFRKRLPMYFWVAGWETCRRMLAFLKD 241 KENRRLALRTRLIRVKAVSKERSAEP 10

Seq. ID No.25

1 gtggccagca gaagtcccca ctccgctgcg ggtggttggc taattcttgg cggctccctt 61 cttgtggtcg gcgtggcgca tccggtagga ctcgccggag gtgacgacga tgctggcgtg 121 gtgcagcagc cgatcgagga tgctggcggc ggtggtgtgc tcgggcagga atcgcccca 181 ttgttcgaag ggccaatgcg aggcgatggc cagggagcgg cgctcgtagc cggcagccac 241 gagccggaac aacagttgag tcccggtgtc gtcgagcggg gcgaagccga tctcgtccaa 301 gatgaccaga teegegegga geagggtgte gatgatettg eegaeggtgt tgteggeeag 361 gccgcggtag aggacctcga tcaggtcggc ggcggtgaag tagcggactt tgaatccggc 421 gtggacggca gcgtgcccgc agccgatgag caggtgactt ttgcccgtac caggtgggcc 481 aatgaccgcc aggttctgtt gtgcccgaat ccattccagg ctcgacaggt agtcgaacgt 541 ggctgcggtg atcgacgatc cggtgacgtc gaacccgtcg agggtcttgg tgaccgggaa 601 ggctgcggcc ttgagacggt tggcggtgtt ggaggcatcg cgggcagcga tctcggcctc 661 aaccaacgte egeaggatet eeteeggtgt eeagegttge gtettggega ettgeaacae 721 ctcggcggcg ttgcggcgca ccgtggccag cttcaaccgc cgcagcgccg cgtcaaggtc 781 ageagecage ggtgeegeeg aggaeggtge caceggettg geageggtgg teatgaggee 841 gtcccgtcgg tggtgttgat cttgtag

Seq. ID No.26

1 VASRSPHSAAGGWLILGGSLLVVGVAHPVG LAGGDDDAGVVQQPIEDAGGGGVLGQESPP L F E G P M R G D G Q G A A L V A G S H E P E Q Q L S P G V 30 V E R G E A D L V Q D D Q I R A E Q G V D D L A D G V V G Q AAVEDLDQVGGGEVADFESGVDGSVPAADE QVTFARTRWANDRQVLLCPNPFQARQVVER G C G D R R S G D V E P V E G L G D R E G C G L E T V G G V GGIAGSDLGLNQRPQDLLRCPALRLGDLQH 35 LGGVAAHRGQLQPPQRRVKVSSQRCRRGRC 271 HRLGSGGHEAVPSVVLIL

- 52 -

Seq. ID No.27

```
1 atgggctgcc tcaaaggtgg tgtcgtcgcc aatgttgttg ttccaacacc ggattatgtg
               61 cgattcgcgt cccactatgg cttcgttccg gacttctgcc acggtgcgga tccgcaatcg
              121 aagggcatcg tggagaacct ctgtggctac gctcaggacg accttgcggt gccgctgctg
              181 accgaagetg egttageegg tgageaggte gacetaegtg eceteaaege ecaggegeaa
5
              241 ctatggtgcg ccgaggtcaa tgccacggtc cactcggaga tctgcgccgt gcccaacgat
              301 cgcttggttg acgagcgcac cgtcttgagg gagctgccct cgctgcggcc gacgatcggc
              361 toggggtogg tgcgccgtaa ggtcgacggc ctctcgtgca tccgttacgg ctcagctcgt
              421 tactoggtgc ctcagoggct cgtcggtgcc accgtggcgg tggtggtcga tcatggcgcc
              481 ctgatcctgt tggaacctgc gaccggtgtg atcgtggccg agcacgagct cgtcagccca
10
              541 ggtgaggtgt ccatcctcga tgaacactac gacggaccca gacccgcacc ctcgcgtggt
              601 cctcgcccga aaacccaagc agagaaacga ttctgcgcat tgggaaccga agcgcagcag
              661 tteetegteg gtgetgetge gateggeaae accegaetga aateegaaet egacattetg
              721 ctcggccttg gcgccgccca cggcgaacag gctttgattg acgcgctgcg ccgggcggtt
              781 gcgtttegec ggtteegege tgeegaegtg egetegatee tggeegeegg egeeggeaee
15
              841 ccacaacccc gccccgccgg cgacgcactc gtgctcgatc tgcccaccgt cgagacccgc
              961 ageoggtgge accettecteg geggeacege tggetgetga cettgaegeg gegetgegge
             1021 ggttgaaget ggccacggtg cgccgcaacg ccgccgaggt gttgcaagtc gccaagacgc
             1081 aacgctggac accggaggag atcctgcgga cgttggttga ggccgagatc gctgcccgcg
20
             1141 atgectecaa cacegocaac egteteaagg cegeageett ceeggteace aagaceeteg
             1201 acgggttcga cgtcaccgga tcgtcgatca ccgcagccac gttcgactac ctgtcgagcc
             1261 tggaatggat tegggeacaa eagaacetgg eggteattgg eecaeetggt aegggeaaaa
             1321 gtcacctgct catcggctgc gggcacgctg ccgtccacgc cggattcaaa gtccgctact
             1381 teacegeege egacetgate gaggteetet acegeggeet ggeegacaac acegteggea
25
             1441 agatcatega caccetgete egegeggate tggtcatett ggacgagate ggettegece
             1501 egetegaega cacegggaet caactgttgt teeggetegt ggetgeegge tacgagegee
             1561 getecetgge categeeteg cattggeet tegaacaatg ggggegatte etgeeegage
             1621 acaccaccge egecageate etegategge tgetgeacca egecageate gtegteacct
              1681 ccggcgagtc ctaccggatg cgccacgccg accacaagaa gggagccgcc aagaattag
30
```

Seq. ID No.28

:0

-

=

н

قت:

4

Į.

1 M G C L K G G V V A N V V V P T P D Y V R F A S H Y G F V P

31 D F C H G A D P Q S K G I V E N L C G Y A Q D D L A V P L L

61 T E A A L A G E Q V D L R A L N A Q A Q L W C A E V N A T V

91 H S E I C A V P N D R L V D E R T V L R E L P S L R P T I G

121 S G S V R R K V D G L S C I R Y G S A R Y S V P Q R L V G A

151 T V A V V V D H G A L I L L E P A T G V I V A E H E L V S P

181 G E V S I L D E H Y D G P R P A P S R G P R P K T Q A E K R

211 F C A L G T E A Q Q F L V G A A A I G N T R L K S E L D I L

40

241 L G L G A A H G E Q A L I D A L R R A V A F R R F R A A D V

271 R S I L A A G A G T P Q P R P A G D A L V L D L P T V E T R

301 S L E A Y K I N T T D G T A S

1 M T T A A K P V A P S S A A P L A A D L D A A L R R L K L A
31 T V R R N A Å E V L Q V A K T Q R W T P E E I L R T L V E A
61 E I A A R D A S N T A N R L K A A A F P V T K T L D G F D V
91 T G S S I T A A T F D Y L S S L E W I R A Q Q N L A V I G P
121 P G T G K S H L L I G C G H A A V H A G F K V R Y F T A A D
151 L I E V L Y R G L A D N T V G K I I D T L L R A D L V I L D
181 E I G F A P L D D T G T Q L L F R L V A A G Y E R R S L A I
211 A S H W P F E Q W G R F L P E H T T A A S I L D R L L H H A

Seq. ID No.30

1 grgacgterg ereegacegt ereggrgata acgaretegt reaacgacet egacgggtrg 61 cagegeaegg tgaaaagtgt gegggegeaa egetaeeggg gaegeatega geacategta 121 ategaeggtg geageggega egaegtggtg geatacetgt eegggtgtga aceaggette 181 gegtattgge agteegagee egaeggeggg eggtaegaeg egatgaacea gggeategeg 15 241 cacgcategg gtgatetgtt gtggttettg cacteegeeg ategttttte egggeeegae 301 gtggtagccc aggccgtgga ggcgctatcc ggcaagggac cggtgtccga attgtggggc 361 ttegggatgg ategtetegt egggetegat egggtgegeg geeegataee ttteageetg 421 cgcaaattee tggeeggeaa geaggttgtt eegeateaag categttett eggateateg 481 ctggtggcca agatcggtgg ctacgacctt gatttcggga tcgccgccga ccaggaattc 20 541 atattgeggg cegegetggt atgegageeg greacgatte ggtgtgtget gtgegagtte 601 gacaccaegg gegteggete geacegggaa ceaagegegg tetteggtga tettegecege 661 atgggcgacc ttcatcgccg ctacccgttc gggggaaggc gaatatcaca tgcctaccta 721 cgcggccggg agttctacgc ctacaacagt cgattctggg aaaacgtctt cacgcgaatg 25 781 tcgaaatag

Seq. ID No.31

1 M T S A P T V S V I T I S F N D L D G L Q R T V K S V R A Q
31 R Y R G R I E H I V I D G G S G D D V V A Y L S G C E P G F
61 A Y W Q S E P D G G R Y D A M N Q G I A H A S G D L L W F L
30 91 H S A D R F S G P D V V A Q A V E A L S G K G P V S E L W G
121 F G M D R L V G L D R V R G P I P F S L R K F L A G K Q V V
151 P H Q A S F F G S S L V A K I G G Y D L D F G I A A D Q E F
181 I L R A A L V C E P V T I R C V L C E F D T T G V G S H R E
211 P S A V F G D L R R M G D L H R R Y P F G G R R I S H A Y L
35 241 R G R E F Y A Y N S R F W E N V F T R M S K



1 gtgaagcgag cgctcatcac cggaatcacc ggccaggacg gctcgtatct cgccgaactg 61 ctgctggcca aggggtatga ggttcacggg ctcatccggc gcgcttcgac gttcaacacc 121 tegeggateg atcaceteta egtegaceeg caccaacegg gegegegget gtttetgeac 181 tatggtgacc tgatcgacgg aacccggttg gtgaccctgc tgagcaccat cgaacccgac 5 241 gaggtgtaca acctggcggc gcagtcacac gtgcgggtga gcttcgacga acccgtgcac 301 accggtgaca ccaccggcat gggatccatg cgactgctgg aagccgttcg gctctctcgg 161 gtgcactgcc gcttctatca ggcgtcctcg tcggagatgt tcggcgcctc gccgccaccg 421 cagaacgage tgacgccgtt ctacccgcgg teaccgtatg gegeegeeaa ggtetatteg 481 tactgggcga cccgcaatta tcgcgaagcg tacggattgt tcgccgttaa cggcatcttg 10 541 ttcaatcacg aatcaccgcg gcgcggtgag acgttcgtga cccgaaagat caccagggcc 601 gtggcacgca tcaaggccgg tatccagtcc gaggtctata tgggcaatct ggatgcggtc 661 cgcgactggg ggtacgcgcc cgaatacgtc gaaggcatgt ggcggatgct gcagaccgac 721 gagcccgacg acttcgtttt ggcgaccggg cgcggtttca ccgtgcgtga gttcgcgcgg 781 gccgcgttcg agcatgccgg tttggactgg cagcagtacg tgaaattcga ccaacgctat 15 841 ctgcggccca ccgaggtgga ttcgctgatc ggcgacgcga ccaaggctgc cgaattgctg 901 ggctggaggg cttcggtgca cactgacgag ttggctcgga tcatggtcga cgcggacatg 961 geggegetgg agtgegaagg caageegtgg ategacaage egatgatege eggeeggaca 1021 tga

20 Seq. ID No.33

1 M K R A L I T G I T G Q D G S Y L A E L L L A K G Y E V H G
31 L I R R A S T F N T S R I D H L Y V D P H Q P G A R L F L H
61 Y G D L I D G T R L V T L L S T I E P D E V Y N L A A Q S H
91 V R V S F D E P V H T G D T T G M G S M R L L E A V R L S R
121 V H C R F Y Q A S S S E M F G A S P P P Q N E L T P F Y P R
151 S P Y G A A K V Y S Y W A T R N Y R E A Y G L F A V N G I L
181 F N H E S P R R G E T F V T R K I T R A V A R I K A G I Q S
211 E V Y M G N L D A V R D W G Y A P E Y V E G M W R M L Q T D
241 E P D D F V L A T G R G F T V R E F A R A A F E H A G L D W
30 271 Q Q Y V K F D Q R Y L R P T E V D S L I G D A T K A A E L L
301 G W R A S V H T D E L A R I M V D A D M A A L E C E G K P W
331 I D K P M I A G R T

Seq. ID No.34

1 atgaggetgg eccgtegege teggaacate ttgegtegea aeggeatega ggtgtegege 61 tactttgccg aactggactg ggaacgcaat ttcttgcgcc aactgcaatc gcatcgggtc 35 181 ggettegegg geegeategt etegttegag eegetgeeeg ggeeetttge egtettgeag 241 cgcagcgcct ccacggaccc gttgtgggaa tgccggcgct gtgcgctggg cgatgtcgat 301 ggaaccatet egateaaegt egeeggeaae gagggegeea geagtteegt ettgeegatg 361 ttgaaacgac atcaggacgc ctttccacca gccaactacg tgggcgccca acgggtgccg 40 421 atacategae tegatteegt ggetgeagae gttetgegge ceaacgatat tgegttettg 481 aagategaeg tteaaggatt egagaageag gtgategegg gtggegatte aaeggtgeae 541 gaccgatgcg tcggcatgca gctcgagctg tctttccagc cgttgtacga gggtggcatg 601 ctcatccgcg aggcgctcga tctcgtggat tcgttgggct ttacgctctc gggattgcaa 661 cccggtttca ccgacccccg caacggtcga atgctgcagg ccgatggcat cttcttccgg 45 721 ggcagcgatt ga

15

20





Seq. ID No.35

1 M R L A R R A R N I L R R N G I E V S R Y F A E L D W E R N
31 F L R Q L Q S H R V S A V L D V G A N S G Q Y A R G L R G A
61 G F A G R I V S F E P L P G P F A V L Q R S A S T D P L W E
91 C R R C A L G D V D G T I S I N V A G N E G A S S S V L P M
121 L K R H Q D A F P P A N Y V G A Q R V P I H R L D S V A A D
151 V L R P N D I A F L K I D V Q G F E K Q V I A G G D S T V H
181 D R C V G M Q L E L S F Q P L Y E G G M L I R E A L D L V D
211 S L G F T L S G L Q P G F T D P R N G R M L Q A D G I F F R

Seq. ID No.36

1 gtgaaatcgt tgaaactcge tegtttcate gegegtageg cegeettega ggtttegege 61 egetattetg agegagacet gaagcaceag tttgtgaage aacteaaate gegtegggta 121 gatgtegttt tegatgtegg egecaactea ggaeaataeg eegeeggeet eegeeggaea 181 geatataagg geegeattgt etegttegaa eegetateeg gaeegtttae gatettggaa 241 ageaaagegt caaeggatee actttgggat tgeeggeage atgegttggg egattetgat 301 ggaaeggtta egateaatat egeaggaaae geeggteaga geagtteegt ettgeeatg 361 etgaaaagte ateagaaege tttteeeeeg geaaaetatg teggtaeeea agaggegtee 421 atacategae ttgatteegt ggegeeagaa tttetaggea tgaaeggtgt egetttete 481 aaggtegaeg tteaaggett tgaaaageag gtgetegeeg ggggeaaate aaceatagat 541 gaeeattgeg teggeatgea tetegtgat teetteetge egttgtaega aggtggeatg 661 eettgttea ttgatgeaaa taatggtega atgttgeagg eegaeggeat etttteeege 721 gaggaegatt ga

25 Seq. ID No.37

1 M K S L K L A R F I A R S A A F E V S R R Y S E R D L K H Q
31 F V K Q L K S R R V D V V F D F T V G A N S G Q Y A A G L R
61 R A A Y K G R I V S F E P L S G P F T I L E S K A S T D P L
91 W D C R Q H A L G D S D G T V T I N I A G N A G Q S S S V L
30 121 P M L K S H Q N A F P P A N Y V G T Q E A S I H R L D S V A
151 P E F L G M N G V A F L K V D V Q G F E K Q V L A G G K S T
181 I D D H C V G M Q L E L S F L P L Y E G G M L I P E A L D L
211 V Y S L G F T L T G L L P C F I D A N N G R M L Q A D G I F

```
1 atggtgcaga cgaaacgata cgccggcttg accgcagcta acacaaagaa agtcgccatg
               61 gccgcaccaa tgttttcgat catcatcccc accttgaacg tggctgcggt attgcctgcc
              121 tgcctcgaca gcatcgcccg tcagacctgc ggtgacttcg agctggtact ggtcgacggc
              181 ggctcgacgg acgaaaccct cgacatcgcc aacattttcg cccccaacct cggcgagcgg
 5
              241 ttgatcattc atcgcgacac cgaccagggc gtctacgacg ccatgaaccg cggcgtggac
              301 ctggccaccg gaacgtggtt gctctttctg ggcgcggacg acagcctgta cgaggctgac
              361 accordgege gggrggeege etteattgge gaacacgage ecagegatet ggratatgge
              421 gacgtgatca tgcgctcaac caatttccgc tggggtggcg ccttcgacct cgaccgtctg
              481 ttgttcaagc gcaacatctg ccatcaggcg atcttctacc gccgcggact cttcggcacc
10
              541 ateggteect acaaceteeg etacegggte etggeegaet gggaetteaa tattegetge
              601 ttttccaacc cagcgctcgt cacccgctac atgcacgtgg tcgttgcaag ctacaacgaa
              661 ttcggcgggc tcagcaatac gatcgtcgad aaggagtttt tgaagcggct gccgatgtcc
              721 acgagactog gcataaggot ggtcatagtt ctggtgcgca ggtggccaaa ggtgatcagc
15
              781 agggccatgg taatgcgcac cgtcatttet tggcggcgcc gacgttag
```

Seq. ID No.39

1 M V Q T K R Y A G L T A A N T K K V A M A A P M F S I I I P

31 T L N V A A V L P A C L D S I A R Q T C G D F E L V L V D G

61 G S T D E T L D I A N I F A P N L G E R L I I H R D T D Q G

91 V Y D A M N R G V D L A T G T W L L F L G A D D S L Y E A D

121 T L A R V A A F I G E H E P S D L V Y G D V I M R S T N F R

151 W G G A F D L D R L L F K R N I C H Q A I F Y R R G L F G T

181 I G P Y N L R Y R V L A D W D F N I R C F S N P A L V T R Y

211 M H V V V A S Y N E F G G L S N T I V D K E F L K R L P M S

25 241 T R L G I R L V I V L V R R W P K V I S R A M V M R T V I S

Seq 40:

GATGCCGTGAGGAGGTAAAGCTGC

Seq 41:

30 GATACGGCTCTTGAATCCTGCACG